

Bowdoin College

Bowdoin Digital Commons

Biology Faculty Publications

Faculty Scholarship and Creative Work

1-1-2019

Mitochondrial genotype influences the response to cold stress in the European green crab, *Carcinus maenas*

Aidan F. Coyle
Bowdoin College

Erin R. Voss
Bowdoin College

Carolyn K. Tepolt
Woods Hole Oceanographic Institution

David B. Carlon
Bowdoin College

Follow this and additional works at: <https://digitalcommons.bowdoin.edu/biology-faculty-publications>

Recommended Citation

Coyle, Aidan F.; Voss, Erin R.; Tepolt, Carolyn K.; and Carlon, David B., "Mitochondrial genotype influences the response to cold stress in the European green crab, *Carcinus maenas*" (2019). *Biology Faculty Publications*. 4.

<https://digitalcommons.bowdoin.edu/biology-faculty-publications/4>

This Article is brought to you for free and open access by the Faculty Scholarship and Creative Work at Bowdoin Digital Commons. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of Bowdoin Digital Commons. For more information, please contact mdoyle@bowdoin.edu, a.sauer@bowdoin.edu.

RESEARCH ARTICLE

Mitochondrial genotype influences the response to cold stress in the European green crab, *Carcinus maenas*

Aidan F. Coyle^{1,*}, Erin R. Voss^{1,‡}, Carolyn K. Tepolt² and David B. Carlon^{1,§}

ABSTRACT

Hybrid zones provide natural experiments in recombination within and between genomes that may have strong effects on organismal fitness. On the East Coast of North America, two distinct lineages of the European green crab (*Carcinus maenas*) have been introduced in the last two centuries. These two lineages with putatively different adaptive properties have hybridized along the coast of the eastern Gulf of Maine, producing new nuclear and mitochondrial combinations that show clinal variation correlated with water temperature. To test the hypothesis that mitochondrial or nuclear genes have effects on thermal tolerance, we first measured the response to cold stress in crabs collected throughout the hybrid zone, then sequenced the mitochondrial *CO1* gene and two nuclear single nucleotide polymorphisms (SNPs) representative of nuclear genetic lineage. Mitochondrial haplotype had a strong association with the ability of crabs to right themselves at 4.5°C that was sex specific: haplotypes originally from northern Europe gave male crabs an advantage while there was no haplotype effect on righting in female crabs. By contrast, the two nuclear SNPs that were significant outliers in a comparison between northern and southern *C. maenas* populations had no effect on righting response at low temperature. These results add *C. maenas* to the shortlist of ectotherms in which mitochondrial variation has been shown to affect thermal tolerance, and suggest that natural selection is shaping the structure of the hybrid zone across the Gulf of Maine. Our limited genomic sampling does not eliminate the strong possibility that mito-nuclear co-adaptation may play a role in the differences in thermal phenotypes documented here. Linkage between mitochondrial genotype and thermal tolerance suggests a role for local adaptation in promoting the spread of invasive populations of *C. maenas* around the world.

KEY WORDS: Cold tolerance, Gulf of Maine, Hybrid zone, Mitochondria, Mito-nuclear interactions, Natural selection, OCLTT model, Oxygen and capacity-limited thermal tolerance

INTRODUCTION

Hybrid zones are geographic regions where distinct lineages or species meet and mate, often over many generations (Harrison, 1993). The genetic novelty maintained in hybrid zones provides

unique opportunities to examine the linkages between genotype and phenotype (Rieseberg et al., 1999). In particular, hybrid zones provide natural experiments to examine how interactions between novel combinations of genes affect physiological and ecological performance. These include interactions between nuclear genes that result from hybridization and backcrossing (Pereira et al., 2014), and interactions across genomes as a result of new cyto-nuclear combinations (Dowling et al., 2008).

Before the Anthropocene, most hybrid zones resulted from shifting range limits driven by Pleistocene climate change that brought formerly non-overlapping lineages or recently evolved species into secondary contact (Hewitt, 2000). More recently, hybrid zones may also be established by repeated biological invasions into the same geographic region from different source populations. Such is the case of the European green crab, *Carcinus maenas* (Linnaeus 1758), which has been introduced into all the major continents of Earth, with the exception of Antarctica, over last two centuries (Darling et al., 2008). On the East Coast of North America, a double invasion has established two distinct genetic lineages across a north–south transect (Roman, 2006). The species was first sighted in coastal Massachusetts in 1817, after which its range slowly expanded north until ~1960 when it appeared to reach its northern range limit near Halifax, Nova Scotia (Carlton and Cohen, 2003). In the early 1980s, this range limit began to move northward, when *C. maenas* was sighted further north and in much colder waters (Roman, 2006). Since 1980, *C. maenas* has expanded northward by roughly 725 km, now spanning the coastal waters of Nova Scotia, Prince Edward Island and a significant portion of Newfoundland (Blakeslee et al., 2010). Analysis of nuclear and mitochondrial DNA (mtDNA) has revealed that this rapid northern expansion was most likely facilitated by a second genetically distinct introduction (Roman, 2006). The first introduction has been traced back to a likely source population in southern or central Europe, while the second introduction consisted of individuals from a northern European source population. Over the last 10 years, the northern lineage has been spreading southward, while the southern lineage has been spreading northward, leading to a region of hybridization and admixture located along the eastern Gulf of Maine and eastern Nova Scotia (Pringle et al., 2011; Darling et al., 2014). The structure of this zone presents a number of interesting experimental opportunities to examine links between genotype and phenotype. For example, mitochondrial haplotypes are spreading more rapidly than nuclear markers, leading to strong mito-nuclear discordance outside the ‘nuclear’ hybrid zone, and novel mito-nuclear combinations within the region of nuclear admixture (Darling et al., 2014; Jeffery et al., 2017). Furthermore, the two lineages are mixing across an ecological seascape that varies in a number of physical and biological factors. Perhaps most importantly, seasonal temperature dynamics vary considerably from Cape Cod to the Gulf of St Lawrence – including a strong gradient in average winter

¹Department of Biology & Schiller Coastal Studies Center, Bowdoin College, Brunswick, ME 04011, USA. ²Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA.

*Present address: Alaska Department of Fish & Game, 351 Research Court, Kodiak, AK 99615, USA. ‡Present address: Museum of Vertebrate Zoology, 3101 Valley Life Sciences Building, University of California Berkeley, Berkeley, CA 94720, USA.

§Author for correspondence (dcarlon@bowdoin.edu)

© A.F.C., 0000-0002-3335-840X; E.R.V., 0000-0003-2662-1190; D.B.C., 0000-0002-5430-4638

temperature that reaches minima of 0–2°C in the Gulf of St Lawrence (Lehnert et al., 2018).

Previous analyses of hybridization dynamics in this complex of closely related populations have largely assumed that mitochondrial variation within and among populations is selectively neutral, and that a number of stochastic demographic factors can explain different rates of migration of genes located in the mitochondria versus the nucleus (Darling et al., 2014). However, to our knowledge there have been no direct studies of how genotypic variation (either mitochondrial or nuclear) might translate into phenotypic variation in this system. There is some indirect evidence that genetic variation may be linked to thermal stress phenotypes from population-level studies. Tepolt and Somero (2014) showed that *C. maenas* collected from different regions within Europe and the USA had distinct responses to heat and cold stress. These same populations showed fairly strong population differentiation in transcriptome-derived single nucleotide polymorphisms (SNPs) across thermal gradients, though sample sizes were too small to test individual-level genotype–phenotype associations conclusively (Tepolt, 2014; Tepolt and Palumbi, 2015). Other studies have found differences in foraging and competitive behavior between *C. maenas* populations from sites with different histories and presumably different genetic backgrounds, leading to the hypothesis that green crabs with ‘northern’ genetic backgrounds may outcompete their southern brethren (Haarr and Rochette, 2012; Rosson et al., 2012). Taken together, this work raises the possibility that thermal stress thresholds in *C. maenas* have genetic components.

In this study, we tested the hypothesis that mitochondrial, nuclear and mito-nuclear interactions can explain differential responses to cold temperature stress, using natural variation in the *C. maenas* hybrid zone in the Gulf of Maine. We chose to focus on mitochondrial variation because the mitochondrial genome codes for a number of key genes that control physiological processes. Variation in mtDNA has been shown to have effects on fitness, often linked to thermal tolerance, in a variety of different animal species. These include flies (Camus et al., 2017), tuna (Dalziel et al., 2006), hares (Ben Slimen et al., 2017) and humans (Balloux et al., 2009). In mammals, variance in the *ATP6* mitochondrial gene has been connected to shifts in temperature tolerance. Mechanistically, *ATP6* variants reduce the coupling efficiency of oxidative phosphorylation, which increases heat production while simultaneously increasing basal metabolic rate (Ballard and Whitlock, 2004). This mechanism results in a trade-off – an individual with an *ATP6* variant has increased cold tolerance at the expense of a higher metabolic rate in more temperate environments – and illustrates how mtDNA can be under selection with thermal tolerance as the selective factor. Furthermore, the mitochondrial genome has a great deal of interaction with the nuclear genome. Mitochondrial proteins, which play an essential role in the oxidative phosphorylation (OXPHOS) system, also interact extensively with nuclear proteins (Burton and Barreto, 2012). The mitochondrial genome is also particularly prone to accumulating deleterious point mutations, and the nuclear genome can take over the function of mitochondrial genes that are knocked out by deleterious substitutions, which may explain the long-term shrinkage of the mitochondrial genome (Rand et al., 2004; Gray, 2011). This tight co-evolution between mitochondrial and nuclear genes can break down in hybrid zones when mtDNA and nuclear DNA are no longer performing complementary roles (Burton and Barreto, 2012). Thus, we expect mito-nuclear interactions to be important in a variety of traits linked to fitness in a broad variety

of systems, and that these interactions are most likely to express themselves in hybrid zones, where new combinations of mitochondrial genomes and nuclear genes are generated.

Our nuclear markers target regions identified in ongoing work on temperature-associated differentiation in cardiac transcriptomes between northern and southern *C. maenas* populations in Europe and the East Coast (Tepolt and Palumbi, 2015). We designed primers for several ‘outlier loci’ identified in that study [loci with high F_{ST} (inbreeding coefficient of the subpopulation compared with the total population) values across these regions] and used PCR primers to sequence and genotype two SNP markers in nuclear regions that showed strong frequency differences between northern and southern Europe. Variation at one of these SNPs is characteristic of southern Europe, while variation at the other SNP is characteristic of northern Europe. Thus, by capitalizing on recombination that is naturally occurring within the *C. maenas* nuclear genome, and between nuclear and mitochondrial genes, we tested whether mitochondrial and selected nuclear genotypes were associated with cold stress phenotypes in wild-caught crabs collected along the hybrid zone on the east coast of North America.

MATERIALS AND METHODS

Collecting sites

To make sure that our experimental assay included genetic variation that was representative of the two ends and middle of the genetic cline, we conducted preliminary sampling and genotyping in Harpswell, ME, USA (N 43°47′27″, W 69°57′34″) and Kent Island, NB, Canada (N 44°33′00″, W 66°45′23″) and found that Harpswell had a high frequency of southern *mt-COI* haplotypes while Kent Island was near the cline center with mixtures of southern and northern haplotypes. To make sure we sampled from the northern end of the cline, we chose Pomquet, NS, Canada (N 45°38′27″, W 61°48′41″) based on previous studies that revealed a high frequency of northern *mt-COI* haplotypes at this site (Roman, 2006). In addition to the 251 crabs collected for the experimental assay, we sequenced 15 crabs from Halifax, NS, Canada (N 44°38′28″, W 63°55′45″) and 22 crabs from Isles of Shoals, ME, USA (N 42°59′23″, W 70°36′58″). These additional samples were used solely to establish geographical clines in our molecular markers and were not used in the experimental assay (Fig. 1). Crabs were collected either by hand in the low intertidal zone or via baited traps placed in the shallow subtidal zone. In Canada, crabs were collected under Fisheries and Oceans Canada Fishing License no. 342021 and 345689. In Maine, crabs were collected under Department of Marine Resources Special License no. ME 2018-24-00.

Cold tolerance assay

The righting response – the time taken for a crab to right itself after being placed on its carapace, ventral-side up – has been shown to be an excellent proxy for other physiological stress variables in a variety of decapods, including *C. maenas* (Cuculescu et al., 1998; Ern et al., 2015). It is also quick to assay, making it easy to test large numbers of crabs. For these reasons, righting response was chosen as the measure of cold tolerance. The cold tolerance assay was conducted at the Schiller Coastal Studies Center in Harpswell, ME, USA. To control for acclimatization, which can have a strong impact on cold tolerance (Tepolt and Somero, 2014), crabs were acclimated in a lab-based flowing seawater system for 3–5 weeks before experimentation at 15.5±0.1°C with a flowrate of 3.79 l min⁻¹. Each tank held crabs from one site at a density of 28–60 crabs per 379 l tank. Tanks were exposed to a 12 h:12 h light:dark cycle with overhead fluorescent light fixtures, and crabs were fed with herring




Fig. 1. Map of *Carcinus maenas* collection sites in the Gulf of Maine. The pie charts depict frequencies of three mitochondrial *CO1* haplogroups at each site: A, red; B, blue; C, black. See Fig. 2 for evolutionary relationships among haplotypes. Sample sizes (n) for mitochondrial genotyping and the number of animals used in the cold stress assay are given below the pie charts for each site.

or shrimp three times a week. To ensure that differential food consumption did not impact behavior, crabs were starved 4–5 days before the start of the testing.

To assay the righting response, crabs were placed in a large plastic tote filled with aerated seawater cooled to 4.5°C in a Percival incubator (Percival Scientific, Perry, IA, USA). For each trial, the individual crabs were subjected to a 5 min cooling period, after which a metal ruler was used to flip the crab onto its back, and the time to right was measured with a stopwatch. All cold tolerance assays were conducted between 07:00 h and 13:00 h on a single day. The time to right was defined as the period between when the animal was flipped on its dorsal side and when the rearmost dactyls maintained contact with the surface of the tote. As the majority of crabs either flipped rapidly within the 2 min trial or failed altogether to right themselves, we used a binary variable for righting response over the continuous variable righting time. After righting response was scored, a variety of morphological parameters – sex, carapace width, number of legs and morph type (red or green) – were recorded for each crab and used as co-variables in statistical models. A single leg of each crab was preserved in 95% ethanol for DNA analysis.

Genotyping

We sequenced a portion of the mitochondrial *CO1* gene (*mt-CO1*) as a marker for mitochondrial haplotype using the primers and PCR conditions published in Roman and Palumbi (2004). The two nuclear markers used targeted coding regions and flanking regulatory regions isolated and sequenced by Tepolt and Palumbi (2015). Three nuclear markers were initially developed from contigs that were identified as outliers in a gene scan between populations located across latitudinal clines in Europe and North America (Dryad repository: <https://doi.org/10.5061/dryad.g8b96/1>). These three initial outlier SNPs were selected as likely deriving from the second *C. maenas* introduction to the east coast. In samples collected in 2011–2012, these SNPs were variable in the Canadian province of Newfoundland and in Norway but fixed for the major allele in Portugal and in the American states of New Jersey and central Maine (Tepolt and Palumbi, 2015). We designed PCR primers to target SNPs within these contigs, Sanger sequenced a set

of samples from collection sites that spanned the Gulf of Maine, and found that all three SNP markers had clinal structures with cline centers located around Penobscot Bay in the Gulf of Maine (Voss, 2016). We could reliably annotate one of these three loci as ubiquitin-conjugating enzyme E2 H or *UBE2H*, and chose it for additional sequencing and tests for association with the cold stress phenotype. The second nuclear marker we used for this study was also identified in transcriptome-based outlier analyses, and was selected as showing a clinal pattern across Europe and North America. In contrast to *UBE2H*, this SNP appears to have been present in the initial introduction to the east coast; in 2011–2012, it was variable in Portugal, New Jersey and Maine but fixed in Newfoundland and Norway (Tepolt and Palumbi, 2015). This SNP was located in a gene annotated as structural maintenance of chromosomes protein 3, or *SMC*. Primer sequences and PCR conditions are listed in Table S1.

DNA extraction was performed with a Qiagen DNEasy Blood & Tissue Kit (Valencia, CA, USA) following the manufacturer's instructions. Amplified products were prepared for sequencing using an exonuclease and shrimp alkaline phosphatase incubation (Exo-SAP) and Sanger sequenced using the GeneWiz sequencing laboratory (South Plainfield, NJ, USA). The resulting chromatograms were edited and aligned using the software package Geneious (Biomatters Ltd, Auckland, New Zealand). The nuclear sequences were scored for SNPs by inspecting the chromatograms for a single versus double peak, the latter indicative of heterozygous status. For the *mt-CO1* gene, five unique haplotypes were identified from all samples. To identify the evolutionary relationships between our sample sequences and those from Europe, reference *mt-CO1* sequences of *C. maenas* within their native range were downloaded from GenBank (accession nos: AY616437.1–AY616445.1, JQ305941.1–JQ305943.1, JQ306002.1–JQ306003.1, KF369118.1) and combined with our sequence data to construct a TCS haplotype network using the software PopART (<http://popart.otago.ac.nz/>).

For the two nuclear markers, we tested for Hardy–Weinberg expectations within loci, and linkage between loci, using Genepop (<http://genepop.curtin.edu.au/>), which implements the estimators of Raymond and Rousset (1995).




Fig. 2. TCS network of the five CO1 haplotypes sampled in *C. maenas* from five sites on the east coast of North America.

Each node represents a unique haplotype, node size is equal to sampling frequency, and hash marks along vertices indicate the number of mutational steps between haplotypes. Pie chart colors correspond to sampling locations indicated in the key. Green nodes illustrate evolutionary relationships of sampled haplotypes with GenBank references, and sampling site and accession number are listed by each node. Three haplogroups (A, B and C) are defined by haplotype similarity and European sampling origin: A, southern Europe; B, northern Europe; and C, pan-European.

Statistical modeling

To analyze the effects of both categorical (sex, mitochondrial haplotype, nuclear genotype, morph color, number of missing limbs) and continuous (carapace width) independent variables on the righting response, we used a multiple logistic regression model implemented by the R-package PerformanceAnalytics (<https://CRAN.R-project.org/package=PerformanceAnalytics>). As explained earlier, the righting response was categorical, with crabs either righting or failing to right within the 2 min trial period. Two models were run. The first included all reproductive, ecological and morphological variables (sex, morph color type, number of legs, sampling location and carapace width) to determine whether any of these variables were associated with the righting response. The results of this model were used to select a smaller set of relevant variables for the final model, which included sex, sampling location and three genetic terms for mitochondrial and nuclear variation. Mitochondrial variation was binned into three haplogroups (A, B and C; see Fig. 2). Nuclear variation was binned into three categories corresponding to the three genotypic states for each biallelic SNP locus.

RESULTS

Genetic polymorphism in *mt-CO1* and tests of Hardy-Weinberg expectations

Five unique *mt-CO1* haplotypes were sampled from the three Canadian and two USA sites and binned into three haplogroups based on evolutionary similarity (Fig. 2). The two sites in Nova

Scotia had a high frequency of two haplotypes closely related or identical to haplotypes commonly found in northern Europe (B1, B2). These two haplotypes were binned within haplogroup B as ‘northern’ haplotypes. The Nova Scotia samples also had a high frequency of a third haplotype related to a haplotype previously observed in Portugal, and believed to have come to the east coast with the second *C. maenas* introduction (Roman, 2006). This is a pan-European haplotype found to comprise roughly 30–40% of Nova Scotia haplotypes in a previous study (Darling et al., 2008), which agrees with the frequency observed in this study. However, as it was not closely related to either of the northern haplotypes (B1 and B2) it was assigned to haplogroup C. The three haplotypes of the B and C haplogroups dominated the two Nova Scotia sites. The central (Kent Island) and two more southern sites (Harpswell and Isles of Shoals) were dominated by the A1 haplotype, identical to a GenBank sample from France (accession no. AY616437), and a defining haplotype of the initial *C. maenas* introduction to the east coast (Roman, 2006). The remaining sampled haplotype (A2) differed by one mutation from A1, and therefore was included in haplogroup A. Translation of these five haplotypes revealed that all mutations were synonymous, yielding identical protein sequences from these partial *mt-CO1* DNA sequences.

Within each of the five sampling sites, we could not reject Hardy-Weinberg expectations within loci for either nuclear marker, and there was no statistical evidence for linkage between *UBE2H* and *SMC* (Table S2). We did observe slightly positive F_{IS} (inbreeding coefficient of an individual relative to the subpopulation) values for




Fig. 3. Geographic clines in molecular markers across the Gulf of Maine. (A) Frequency of the five mitochondrial haplotypes from Fig. 2. (B) Frequency of two nuclear single nucleotide polymorphism (SNP) markers. Sample sizes are given above data points.

UBE2H among all sites, suggesting mild heterozygote deficits at this locus, but these values did not depart from null Hardy–Weinberg expectations.

Clines in mitochondrial and nuclear markers

For *mt-COI*, we observed a strong cline in the frequency of haplotypes (Figs 1 and 3A). The A haplotypes dominated the two southern sites (Isle of Shoals and Harpswell) and declined in frequency beyond Kent Island at the two sites in Nova Scotia. In fact, this pattern was dominated by the A1 haplotype, which comprised 67%, 84%, 70%, 13% and 7% of the samples, moving from south to north, respectively. Conversely, the B1, B2 and C haplotypes increased in frequency at the two sites in Nova Scotia, so that ~75% of the samples carried one of these three haplotypes. Interestingly, we sampled a few ‘northern’ B1 and C haplotypes at Isle of Shoals, a southern site that is offshore; however, the sample size was too small ($n=6$) for more precise estimates of haplotype frequency.

The two nuclear loci showed contrasting clinal patterns (Fig. 3B). The *UBE2H* A nucleotide, likely derived from the second introduction from a northern European source, increased in frequency moving from south to north, while the *SMC* T nucleotide, likely present in the original introduction from a south-central European source, reached

the highest frequency on Kent Island, but decreased at both the southern and northern ends of the distribution.

Multiple logistic regression models of the righting response under cold stress

In our model, which included only ecological, morphological and sexual variables, only site had a significant effect on the righting response ($P=0.0046$), while sex approached marginal significance ($P=0.149$; Table 1). We therefore included both of these terms in the final model, which also had terms for variation at *mt-COI* and

Table 1. The results of a multiple regression logistic model of five variables on the righting response

Factor	Deviance	AIC	LRT	Pr(>Chi)
Sex	195.10	201.27	2.169	0.149
Morph type	197.27	202.96	0.144	0.701
Missing legs	194.96	202.99	0.111	0.739
Carapace width	195.09	203.09	0.012	0.913
Site	203.12	207.12	8.015	0.0046

AIC, Akaike's information criterion; LRT, likelihood ratio test; Pr(>Chi), probability that the chi-square value is significant. Gray shading highlights terms with the largest statistical effects.

Table 2. The results of a multiple regression logistic model that included terms for site, sex, SNP variation at two nuclear loci (*UBE2H* and *SMC*) and *mt-CO1* haplotype variation

Factor	Deviance	AIC	LRT	Pr(>Chi)
Site	156.73	164.73	0.052	0.819
Sex	160.56	164.56	3.782	0.052
<i>UBE2H</i>	156.46	164.46	0.327	0.573
<i>SMC</i>	156.76	164.76	0.0207	0.885
<i>mt-CO1</i>	165.81	169.81	9.0334	0.00265

SNP, single nucleotide polymorphism. The three *mt-CO1* haplogroups are A, B and C as in Fig. 2. Gray shading highlights terms with the largest statistical effects.

nuclear markers. In this second model, sex was marginally significant ($P=0.052$), site was not significant ($P=0.819$) and *mt-CO1* haplogroup was highly significant ($P=0.003$; Table 2). Somewhat surprisingly, variation at either nuclear marker had no effect on righting response in this model ($P\geq 0.573$). The effects of sex and *mt-CO1* terms are graphically presented in Fig. 4. In males, the northern (B) and pan-European (C) haplotypes increased the probability of righting at 4.5°C by ~20% compared with the southern haplotypes (A) (Fig. 4A). In fact, every single male with a northern or pan-European haplotype was able to successfully right itself at 4.5°C. While the righting response in females did not appear to be affected by *mt-CO1* haplogroup, sample sizes were smaller compared with those of males, and were therefore unlikely to uncover small effect sizes, if they indeed exist. The effects of sex and *mt-CO1* haplogroup were consistent across the three sites. While sample sizes of some haplogroup × sex combinations were necessarily small because of the clinal structure of mitochondrial variation, males that carried northern or pan-European haplotypes (B,C) benefited from a ~0.20 increase in the probability of flipping at 4.5°C over males that carried southern haplotypes (A) at both Kent Island and Pomquet (Fig. 4B). Moreover, the effect of carrying an A haplotype was very similar across all three sites: males and females had a ~0.75 probability of flipping at 4.5°C, regardless of site of collection.

DISCUSSION

Connecting genotype to phenotype

Our results indicate a strong and sex-specific effect of mitochondrial haplotype on the ability of green crabs to tolerate cold stress. In our logistic regression model that included terms for site, sex and genetic factors, mitochondrial haplotype was highly significant and had a large effect on the righting response under cold stress (Table 2, Fig. 4). At each site, males carrying B or C haplotypes had a ~20% greater probability of righting at 4.5°C than did their counterparts carrying A haplotypes. It appears that differences in mitochondrial genomes play an important role in differences between sites in terms of cold tolerance: in the model without genetic factors, site of origin is highly significant, but when both site and haplogroup are included, site becomes non-significant. We note that our results do not suggest that the mitochondrial *COI* gene specifically plays a role in cold tolerance. All five haplotypes yielded an identical protein sequence, and as mitochondrial genes do not undergo recombination and the entire mitochondrial genome is inherited as a single haploid locus, our mitochondrial marker is linked to mutations throughout the mitochondrial genome that may affect physiological performance.

Nearshore winter temperatures at the Pomquet site were much colder (<0°C) than those at either Kent Island or Harpswell (~5°C) (Fig. 5). The Pomquet winter temperatures are well below our 4.5°C

assay temperature, and suggest there could be strong selection for cold-tolerant phenotypes along the coast of Nova Scotia and northward. The fact that male cold tolerance was connected to mtDNA haplotype while female cold tolerance was not presents an intriguing evolutionary paradox. Because the mitochondrial genome is typically uniparentally inherited through females, mutations arising in the mitochondrial genome that affect males, but not females, are ‘invisible’ to natural selection (Beekman et al., 2014). Two caveats to this general result are situations where there is either high inbreeding or opportunities for kin selection (Wade and Brandvain, 2009). In *C. maenas*, where the potential for long-distance larval dispersal will likely preclude any substantial inbreeding or kin selection, we see two potential solutions to this paradox. The first posits strong mito-nuclear co-evolution in response to spatially varying temperature stress. Mitochondrial genes are either involved in mtDNA translation or produce proteins needed for the electron transport chain (Ballard and Whitlock, 2004). However, while there are fewer than 40 mitochondrial genes, there are roughly 1500 nuclear genes involved in OXPHOS (Gershoni et al., 2009). Thus, OXPHOS is likely to be strongly dependent on mito-nuclear interactions (Wolff et al., 2014). These protein interactions typically rely on a ‘lock-and-key’ principle, in which the nuclear and mitochondrial proteins must match each other. Support for the general idea that the evolution of stress tolerance requires substantial co-evolution between nuclear and mitochondrial loci comes from a handful of experimental insect studies that have crossed mtDNA variants into divergent nuclear backgrounds (Arnqvist et al., 2010; Koevoets et al., 2012; Hoekstra et al., 2013). These studies have shown that metabolic and life-history traits in F2 hybrids have increased temperature sensitivity when compared with non-recombinant parental lines. In contrast, F2 hybrids between divergent populations of the marine copepod *Tigriopus californicus* have shown patterns of transgressive segregation and novel thermal phenotypes that are inconsistent with mito-nuclear co-evolution (Willett, 2010; Pereira et al., 2014). A second mechanism that would facilitate selection on male traits caused by mtDNA variation are pleiotropic interactions between male and female traits that affect fitness. Camus et al. (2015) have shown strong, antagonistic pleiotropic effects of mtDNA mutations on male fertility and female longevity in *Drosophila melanogaster*, illustrating the importance of documenting the role of mtDNA variation in a broad array of male and female traits to understand how selective gradients are shaping mitochondrial and nuclear variation.

As discussed in the Introduction, there are a handful of other examples of species where mitochondrial variation plays an important role in thermal phenotype. While the majority of these examples are mammalian, there is also accumulating correlative evidence from ectotherms (including insects and fishes) that shows clinal variation in mitochondrial haplotypes maps onto the thermal environment, or that molecular signatures of natural selection are accelerated in comparisons between populations living in different thermal environments (reviewed in Camus et al., 2017). To our knowledge, our study is the first to establish a direct link between mitochondrial variation and temperature tolerance in a marine ectotherm. Copepods are a likely second candidate. Harada et al. (2019) have shown that variation in mitochondrial function can explain differences in tolerance to acute heat stress among populations distributed across a temperature gradient spanning coastal California in the intertidal copepod *Tigriopus californicus*. While mtDNA was not sequenced in this study, these same populations have previously been shown to have between 9.5% and 26.5% sequence divergence among populations. Given that




Fig. 4. Righting response of male and female crabs at 4.5°C. (A) Data from all three sites combined, binned by *mt-CO1* haplogroup. (B) Data for each individual site, binned by *mt-CO1* haplogroup. Sites are vertically arranged from south (Harpowell) to north (Pomquet). The sample size (*n*) of each sex × *mt-CO1* haplogroup combination is given at the base of each bar; *n*=0 indicates that the experimental population did not include individuals of that sex and *mt-CO1* haplogroup. See Table 2 for a statistical model of the effects of sex and *mt-CO1* haplogroup.

temperature tolerance varies among populations and closely related species that partition distinct thermal environments in the intertidal zone (Somero, 2002), we predict a broader role for mitochondrial variation in thermal adaptation in the marine realm.

A genotypic effect on phenotype was not seen for the two nuclear genes chosen in this study, indicating that these particular loci are not linked to genes involved in the righting response in the cold or that our sample sizes were not large enough to detect smaller effect sizes. However, both nuclear markers were outliers in analyses of transcriptomic variation between northern and southern sites, and

SMC was specifically associated with site temperature (Tepolt and Palumbi, 2015). An independent restriction site-associated DNA sequencing (RAD-seq)-based study has also uncovered evidence for potential selection in response to winter temperatures in east coast *C. maenas* populations (Jeffery et al., 2018). Given this, we suggest that future studies should target a wider range of nuclear markers and physiological tests to more fully explore the role of the nuclear genome in setting thermal limits in the system, particularly in the context of mitochondrial lineage-associated differences in cold tolerance.




Fig. 5. Monthly sea surface temperature (SST) data at each tested site for the years 2012–2016. Solid line indicates mean SST for the month across all years, while the gray shading indicates the maximum and minimum temperatures recorded during that period. Data represent the closest 0.25 deg grid location to each site, and were derived from NOAA's OI SST V2 High Resolution Dataset provided by the NOAA/OAR/ESRL PSD (Boulder, CO, USA) from their website: <https://www.esrl.noaa.gov/psd/> (Reynolds et al., 2007).

The main mechanistic hypothesis for how the mitochondrial genome impacts cold tolerance in ectotherms is oxygen and capacity-limited thermal tolerance (OCLTT) (Pörtner and Knust, 2007; Ballard and Melvin, 2010; Ern et al., 2015). According to this hypothesis, cold-adapted ectotherms have a higher mitochondrial capacity but also an increased metabolic rate. Increased cold tolerance is a trade-off, with lower heat tolerance as its result. Direct evidence for a OCLTT trade-off comes from a mtDNA cline in Australian *Drosophila melanogaster*, where laboratory populations generated from isofemale lines sampled from the tropics (Queensland) carried mitochondrial haplotypes with greater heat tolerance and less cold tolerance than mitochondrial haplotypes originating from the temperate location (Brisbane), which had greater cold tolerance and less heat tolerance (Camus et al., 2017). There is some support for a similar OCLTT trade-off between heat and cold tolerance in *C. maenas* populations sampled from different geographical regions (Tepolt and Somero, 2014). Within their native range and the east coast of North America, northern populations dominated by B and C haplogroups are more cold tolerant than southern populations dominated by the A haplogroup (Roman and Palumbi, 2004; Tepolt and Somero, 2014). In turn, the southern populations are more heat tolerant than the northern populations. This apparent trade-off between heat and cold tolerance provides preliminary evidence that mitochondrial adaptation in *C. maenas* may follow the OCLTT model of thermal tolerance. However, the strongest test of the model and its predicted trade-off will be to experimentally subject crabs carrying different mitochondrial haplotypes to both cold and heat stress.

Both of our nuclear markers differ in their clinal patterns from *COI*. Previous sampling revealed that *UBE2H* variation, which likely derives from the second *C. maenas* introduction, changed in allele frequency further south than *COI*, near Penobscot Bay (Voss, 2016) and that the frequency of the T nucleotide does not increase to >0.50 at our two sites in Nova Scotia. In contrast, the *SMC* T nucleotide, deriving from the first *C. maenas* introduction, reaches its highest frequency of 0.33 at Kent Island, but declines in frequency at the more southern and northern sites. Multiple genome-wide studies have shown complex patterns of genomic admixture along the east coast. Initial work with microsatellites in 2007 suggested that mitochondrial DNA was introgressing faster both up and down the coast than nuclear DNA, with 'southern'

haplotypes found with 'northern' nuclear backgrounds outside of the primary introgression zone, and vice versa ('northern' haplotypes in 'southern' backgrounds) (Darling et al., 2014). In 2011, transcriptome sequencing showed extensive admixture in the recent Newfoundland expansion, which is believed to derive from an admixed Nova Scotia population, with no evidence for any nuclear introgression in 'southern' populations in New Jersey and central Maine (Blakeslee et al., 2010; Tepolt and Palumbi, 2015). A geographically broad RAD-seq-based study showed extensive nuclear introgression in southeastern Nova Scotia and southeastern Newfoundland, with most other east coast populations composed primarily or exclusively of animals from a primarily northern or southern nuclear background (Jeffery et al., 2017). Given both this prior work and the nuclear patterns we see here, it is likely that our Harpswell crabs had primarily southern nuclear backgrounds; the Kent Island crabs had admixed nuclear backgrounds; and the Pomquet crabs had northern backgrounds. Thus, the association we observed between cold tolerance and mitochondrial lineage holds true across very different nuclear genetic backgrounds (Fig. 4B). While these mito-nuclear patterns might at first appear to reject the hypothesis of tightly coupled mito-nuclear interactions, to our knowledge none of the previous studies have explicitly tested for linkage between mitochondrial haplotypes and large panels of nuclear SNPs, and it seems likely that these associations would be overwhelmed by the dominant signal from non-linked markers. Future work in extensively admixed populations in southeastern Nova Scotia and Newfoundland that targets genes involved in OXPHOS will help to more fully explore the role of the nuclear genome in mediating the observed effect of mitochondrial lineage on cold tolerance.

The *C. maenas* hybrid zone presents an interesting system in which to study the role of sexual selection in shaping a variety of traits. Sexual dimorphism in morphology (males reach much larger sizes than females) and behavior is thought to result from male-male conflicts and sexual selection (Berrill and Arsenault, 1982). In addition, there is strong intra-sexual selection for male aggressive behavior to guard vulnerable females from other males before and after copulation, which may interact with natural selection to drive sexual dimorphism in cold tolerance. Any energetic advantage of substitutions in the mitochondrial genome, coupled with complementary nuclear variation, might amplify this behavioral

difference between sexes. Mito-nuclear co-evolution will facilitate inter-sexual selection because it requires reliable matching of the mitochondrial genes with nuclear genes (Hill, 2015). Under this paradigm, female preference will evolve in concert with male signals that increase mitochondrial–nuclear matching. Areas where there is extensive natural mixing of mtDNA haplotypes and nuclear backgrounds provide an interesting field test of this hypothesis. While the genetic basis of sex determination in *C. maenas* is unknown, studies in a few other brachyuran decapods have documented both XY (in *Charybdis feriata*, also a portunid) and ZW (in *Erochier sinensis*, a distantly related varunid) systems (reviewed by Chandler et al., 2018). An intriguing possibility emerges in light of the findings that nuclear-encoded genes that function in the mitochondria (N-mt genes) are under-represented on the X chromosome and over-represented on autosomes in XY systems (Drown et al., 2012), but that N-mt genes whose expression has been shown to be sensitive to mtDNA variation are over-represented on the Y chromosome of *D. melanogaster* (Rogell et al., 2014). These latter genes on the Y chromosome are predicted to be responding to mito-nuclear antagonisms that result from maternal inheritance, by compensating for male-harming mtDNA mutations. Could we have experimentally uncovered an ecologically important male phenotype that has roots in sexually antagonistic evolution? Clearly, sex determination, the location of N-mt genes in *C. maenas* and their role in variable cold tolerance are all of considerable interest in this hybrid zone.

Mitochondrial genes as neutral markers?

Previous studies of the dynamics of the green crab hybrid zone have focused on a variety of hypotheses that assume mitochondrial *COI* is a neutral proxy for dispersal and is not under selection (Roman, 2006; Pringle et al., 2011; Darling et al., 2014). More broadly, the field of phylogeography was founded on studies that relied on mitochondrial variation and the assumption of effective neutrality (Avice, 2004). We have established a clear link between mitochondrial haplotype and male cold tolerance in *C. maenas*, which suggests that differences in cline structure between mitochondrial markers and other, putatively neutral, nuclear markers (e.g. microsatellites and SNPs; Darling et al., 2014; Jeffery et al., 2017) are likely related to a shifting balance between selection and stochastic demographic processes that is marker dependent. With regards to gradients in temperature in the Gulf of Maine and the Canadian Maritimes, the influence of the cold Labrador current is particularly strong as it moves southward along the coast of Nova Scotia and into the upper reaches of the Gulf of Maine, essentially splitting the eastern and western Gulf into two different thermal regimes (Pettigrew et al., 2005). Thus, the thermal transition from the stable and colder waters of the eastern Gulf of Maine to the seasonally warmer temperatures in the western Gulf of Maine appears to play a key role in the structure and persistence of the mitochondrial cline, and we predict similar clines in nuclear loci that play key roles in mitochondrial functions.

Further, the functional differences among mtDNA haplotypes imply that adaptation has played a key role in the limitation of the original *C. maenas* invasion to the Gulf of Maine, with the species' expansion into the Canadian Maritime provinces coming only after the introduction of presumably cold-adapted genetic variants (Roman, 2006). Assuming mtDNA differences and their likely co-adapted nuclear-encoded mitochondrial genes represent standing genetic variation originally shaped by the physical differences between northern and southern Europe in its native

range over the last 500,000 years (Roman and Palumbi, 2004; Tepolt and Somero, 2014; Jeffery et al., 2018), then the success of *C. maenas* in North America and elsewhere may be related to pre-adaptation to specific thermal environments. Observational studies have suggested a role for mitochondrial preadaptation in invasion success in other systems, notably the bryozoan *Watersipora subtorquata* (Mackie et al., 2012), but ours presents the first mechanistic support for this possibility. Predicting the success of new invasions may therefore depend on the invading genotypes as well as their interactions with the native community. Furthermore, such preadaptation would have important implications for invasion management, suggesting that there may be advantages to preventing further introductions of invasive species that have already become established in a new region.

Conclusions

Our finding of an association between mtDNA haplotypes and thermal tolerance in invasive green crabs may help to explain the discordance between mtDNA and presumably neutral nuclear markers along the northeastern coast of the USA. The increase in frequency of cold-adapted mitochondrial haplotypes along the coast of Nova Scotia and within the Gulf of St Lawrence coincides with a shift towards greater seasonal extremes in winter temperatures. This finding adds to a growing number of marine and terrestrial studies indicating an important role for mtDNA variation in shaping organismal responses to temperature stress, but also suggests complexity in the genomic architecture of stress-related phenotypes. The male-specific effect of mtDNA variation we observed can be viewed as an evolutionary paradox, whose resolution may reside in mito-nuclear co-adaptation to thermal stress, or in pleiotropic effects of mtDNA variation on male and female traits. With the advent of a rich and increasingly affordable array of genomic tools for natural systems (Sherman et al., 2016), *C. maenas* is an excellent model with which to explore the genomic drivers of a wide variety of physiological, population and ecosystem processes in a super-invader that is profoundly changing benthic seascapes throughout the world. This work highlights the importance of understanding the genetic makeup of invading populations and how these invading genotypes affect ecological phenotypes when predicting the success and ecological outcome of invasions.

Acknowledgements

We would like to thank T. Suskiewicz and L. Johnson for help collecting crabs from Halifax, NS, and Robin Seeley for collections from the Isle of Shoals. We thank Timothy Fuller for designing and testing the SMC primers. We thank Mark Murray for facilitating a productive stay on Kent Island, and Nick Keeney for assistance with animal care at the Schiller Coastal Studies Center. This is publication no. 5 from the Bowdoin Marine Laboratory.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: A.F.C.; Methodology: A.F.C., D.B.C.; Formal analysis: A.F.C., D.B.C.; Investigation: A.F.C., E.R.V., C.K.T., D.B.C.; Resources: C.K.T., D.B.C.; Writing - original draft: A.F.C., D.B.C.; Writing - review & editing: A.F.C., E.R.V., C.K.T., D.B.C.; Visualization: A.F.C.; Supervision: D.B.C.; Funding acquisition: D.B.C.

Funding

A.F.C. was supported by a Grua/O'Connell Research Award from Bowdoin College, and a grant from the Quahog Bay Conservancy. E.R.V. was supported by a Freedman Fellowship for Coastal & Environmental Studies and by a Phocas Family Fellowship from Bowdoin College. D.B.C. was supported by startup funds from Bowdoin College. C.K.T. was supported by The James S. Cole and Cecily C. Selby Endowed Fund in Support of Scientific Staff.

Data availability

The mitochondrial sequence data is available from GenBank under accession nos: MK634747–MK634997. The nuclear sequence data and data from the cold tolerance assay are available from Dryad (Coyle et al., 2019): <https://doi.org/10.5061/dryad.cq91j7v>

Supplementary information

Supplementary information available online at <http://jeb.biologists.org/lookup/doi/10.1242/jeb.203521.supplemental>

References

- Arnqvist, G., Dowling, D. K., Eady, P., Gay, L., Tregenza, T., Tuda, M. and Hosken, D. J. (2010). Genetic architecture of metabolic rate: environment specific epistasis between mitochondrial and nuclear genes in an insect. *Evolution* **64**, 3354–3363. doi:10.1111/j.1558-5646.2010.01135.x
- Avise, J. C. (2004). *Molecular Markers, Natural History and Evolution*, 2nd edn. Basingstoke, UK: Springer Nature.
- Ballard, J. W. O. and Whitlock, M. C. (2004). The incomplete natural history of mitochondria. *Mol. Ecol.* **13**, 729–744. doi:10.1046/j.1365-294X.2003.02063.x
- Ballard, J. W. O. and Melvin, R. G. (2010). Linking the mitochondrial genotype to the organismal phenotype. *Mol. Ecol.* **19**, 1523–1539. doi:10.1111/j.1365-294X.2010.04594.x
- Balloux, F., Handley, L.-J. L., Jombart, T., Liu, H. and Manica, A. (2009). Climate shaped the worldwide distribution of human mitochondrial DNA sequence variation. *Proc. R. Soc. B Biol. Sci.* **276**, 3447–3455. doi:10.1098/rspb.2009.0752
- Beekman, M., Dowling, D. K. and Aanen, D. K. (2014). The costs of being male: are there sex-specific effects of uniparental mitochondrial inheritance. *Philos. T. R. Soc. B.* **369**, 20130440. doi:10.1098/rstb.2013.0440
- Ben Slimen, H., Schaschl, H., Knauer, F. and Suchentrunk, F. (2017). Selection on the mitochondrial ATP synthase 6 and the NADH dehydrogenase 2 genes in hares (*Lepus capensis* L., 1758) from a steep ecological gradient in North Africa. *BMC Evol. Biol.* **17**, 46. doi:10.1186/s12862-017-0896-0
- Berrill, M. and Arsenaault, M. (1982). Mating behavior of the green shore crab *Carcinus maenas*. *Bull. Mar. Sci.* **32**, 632–638.
- Blakeslee, A. M. H., McKenzie, C. H., Darling, J. A., Byers, J. E., Pringle, J. M. and Roman, J. (2010). A hitchhiker's guide to the Maritimes: anthropogenic transport facilitates long distance dispersal of an invasive marine crab to Newfoundland. *Divers. Distrib.* **16**, 879–891. doi:10.1111/j.1472-4642.2010.00703.x
- Burton, R. S. and Barreto, F. S. (2012). A disproportionate role for mt DNA in Dobzhansky–Muller incompatibilities. *Mol. Ecol.* **21**, 4942–4957. doi:10.1111/mec.12006
- Camus, M. F., Wolf, J. B. W., Morrow, E. H. and Dowling, D. K. (2015). Single nucleotides in the mtDNA sequence modify mitochondrial molecular function and are associated with sex-specific effects on fertility and aging. *Curr. Biol.* **25**, 2717–2722. doi:10.1016/j.cub.2015.09.012
- Camus, M. F., Wolff, J. N., Sgrò, C. M. and Dowling, D. K. (2017). Experimental support that natural selection has shaped the latitudinal distribution of mitochondrial haplotypes in Australian *Drosophila melanogaster*. *Mol. Biol. Evol.* **34**, 2600–2612. doi:10.1093/molbev/msx184
- Carlton, J. T. and Cohen, A. N. (2003). Episodic global dispersal in shallow water marine organisms: the case history of the European shore crabs *Carcinus maenas* and *C. aestuarii*. *J. Biogeogr.* **30**, 1809–1820. doi:10.1111/j.1365-2699.2003.00962.x
- Chandler, J. C., Elizur, A. and Ventura, T. (2018). The decapod researcher's guide to the galaxy of sex determination. *Hydrobiologia* **825**, 61–80. doi:10.1007/s10750-017-3452-4
- Coyle, A. F., Voss, E. R., Tepolt, C. K. and Carlton, D. B. (2019). Data from: Mitochondrial genotype influences the response to cold stress in the European green crab *Carcinus maenas*. *Dryad Digital Repository*. <https://doi.org/10.5061/dryad.cq91j7v>
- Cuculescu, M., Hyde, D. and Bowler, K. (1998). Thermal tolerance of two species of marine crab, *Cancer pagurus* and *Carcinus maenas*. *J. Therm. Biol.* **23**, 107–110. doi:10.1016/S0306-4565(98)00008-4
- Dalziel, A. C., Moyes, C. D., Fredriksson, E. and Loughheed, S. C. (2006). Molecular evolution of cytochrome c oxidase in high-performance fish (Teleostei: Scombroidei). *J. Mol. Evol.* **62**, 319–331. doi:10.1007/s00239-005-0110-7
- Darling, J. A., Bagley, M. J., Roman, J. O. E., Tepolt, C. K. and Geller, J. B. (2008). Genetic patterns across multiple introductions of the globally invasive crab genus *Carcinus*. *Mol. Ecol.* **17**, 4992–5007. doi:10.1111/j.1365-294X.2008.03978.x
- Darling, J. A., Tsai, Y.-H. E., Blakeslee, A. M. H. and Roman, J. (2014). Are genes faster than crabs? Mitochondrial introgression exceeds larval dispersal during population expansion of the invasive crab *Carcinus maenas*. *Roy. Soc. Open Sci.* **1**, 140202. doi:10.1098/rsos.140202
- Dowling, D. K., Friberg, U. and Lindell, J. (2008). Evolutionary implications of non-neutral mitochondrial genetic variation. *Trends Ecol. Evol.* **23**, 546–554. doi:10.1016/j.tree.2008.05.011
- Drown, D. M., Preuss, K. M. and Wade, M. J. (2012). Evidence of a paucity of genes that interact with the mitochondrion on the X in mammals. *Genome Biol. Evol.* **4**, 875–880. doi:10.1093/gbe/evs064
- Ern, R., Huong, D. T. T., Phuong, N. T., Madsen, P. T., Wang, T. and Bayley, M. (2015). Some like it hot: thermal tolerance and oxygen supply capacity in two eurythermal crustaceans. *Sci. Rep.* **5**, 10743. doi:10.1038/srep10743
- Gershoni, M., Templeton, A. R. and Mishmar, D. (2009). Mitochondrial bioenergetics as a major motive force of speciation. *BioEssays* **31**, 642–650. doi:10.1002/bies.200800139
- Gray, M. W. (2011). The incredible shrinking organelle. *EMBO Rep.* **12**, 873–873. doi:10.1038/embor.2011.168
- Haarr, M. L. and Rochette, R. (2012). The effect of geographic origin on interactions between adult invasive green crabs *Carcinus maenas* and juvenile American lobsters *Homarus americanus* in Atlantic Canada. *J. Exp. Mar. Biol. Ecol.* **422**, 88–100. doi:10.1016/j.jembe.2012.04.016
- Harada, A. E., Healy, T. M. and Burton, R. S. (2019). Variation in thermal tolerance and its relationship to mitochondrial function across populations of *Tigriopus californicus*. *Front. Physiol.* **10**, 213. doi:10.3389/fphys.2019.00213
- Harrison, R. G. (1993). *Hybrid Zones and the Evolutionary Process*. New York: Oxford University Press.
- Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913. doi:10.1038/35016000
- Hill, G. E. (2015). Mitonuclear ecology. *Mol. Biol. Evol.* **32**, 1917–1927. doi:10.1093/molbev/msv104
- Hoekstra, L. A., Siddiq, M. A. and Montooth, K. L. (2013). Pleiotropic effects of a mitochondrial–nuclear incompatibility depend upon the accelerating effect of temperature in *Drosophila*. *Genetics* **195**, 1129–1139. doi:10.1534/genetics.113.154914
- Jeffery, N. W., DiBacco, C., Wringe, B. F., Stanley, R. R. E., Hamilton, L. C., Ravindran, P. N. and Bradbury, I. R. (2017). Genomic evidence of hybridization between two independent invasions of European green crab (*Carcinus maenas*) in the Northwest Atlantic. *Heredity* **119**, 154. doi:10.1038/hdy.2017.22
- Jeffery, N. W., Bradbury, I. R., Stanley, R. R. E., Wringe, B. F., Van Wyngaarden, M., Lowen, J. B., McKenzie, C. H., Matheson, K., Sargent, P. S. and DiBacco, C. (2018). Genomewide evidence of environmentally mediated secondary contact of European green crab (*Carcinus maenas*) lineages in eastern North America. *Evol. Appl.* **11**, 869–882. doi:10.1111/eva.12601
- Koevoets, T., Van De Zande, L. and Beukeboom, L. W. (2012). Temperature stress increases hybrid incompatibilities in the parasitic wasp genus *Nasonia*. *J. Evol. Biol.* **25**, 304–316. doi:10.1111/j.1420-9101.2011.02424.x
- Lehnert, S. J., DiBacco, C., Jeffery, N. W., Blakeslee, A. M. H., Isaksson, J., Roman, J., Wringe, B. F., Stanley, R. R. E., Matheson, K. and McKenzie, C. H. (2018). Temporal dynamics of genetic clines of invasive European green crab (*Carcinus maenas*) in eastern North America. *Evol. Appl.* **11**, 1656–1670. doi:10.1111/eva.12657
- Mackie, J. A., Darling, J. A. and Geller, J. B. (2012). Ecology of cryptic invasions: latitudinal segregation among Watersipora (Bryozoa) species. *Sci. Rep.* **2**, 871. doi:10.1038/srep00871
- Pereira, R. J., Barreto, F. S. and Burton, R. S. (2014). Ecological novelty by hybridization: experimental evidence for increased thermal tolerance by transgressive segregation in *Tigriopus californicus*. *Evolution* **68**, 204–215. doi:10.1111/evo.12254
- Pettigrew, N. R., Churchill, J. H., Janzen, C. D., Mangum, L. J., Signell, R. P., Thomas, A. C., Townsend, D. W., Wallinga, J. P. and Xue, H. (2005). The kinematic and hydrographic structure of the gulf of maine coastal current. *Deep Sea Res. II* **52**, 2369–2391. doi:10.1016/j.dsr2.2005.06.033
- Pörtner, H. O. and Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* **315**, 95–97. doi:10.1126/science.1135471
- Pringle, J. M., Blakeslee, A. M. H., Byers, J. E. and Roman, J. (2011). Asymmetric dispersal allows an upstream region to control population structure throughout a species range. *Proc. Nat. Acad. Sci. USA* **108**, 15288–15293. doi:10.1073/pnas.1100473108
- Rand, D. M., Haney, R. A. and Fry, A. J. (2004). Cytonuclear coevolution: the genomics of cooperation. *Trends Ecol. Evol.* **19**, 645–653. doi:10.1016/j.tree.2004.10.003
- Raymond, M. and Rousset, F. (1995). An exact test for population differentiation. *Evolution* **49**, 1280–1283. doi:10.1111/j.1558-5646.1995.tb04456.x
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S. and Schlax, M. G. (2007). Daily high-resolution-blended analyses for sea surface temperature. *J. Clim.* **20**, 5473–5496. doi:10.1175/2007JCLI1824.1
- Rieseberg, L. H., Whitton, J. and Gardner, K. (1999). Hybrid zones and the genetic architecture of a barrier to gene flow between two sunflower species. *Genetics* **152**, 713–727.
- Rogell, B., Dean, R., Lemos, B. and Dowling, D. K. (2014). Mito-nuclear interactions as drivers of gene movement on and off the X-chromosome. *BMC Genom.* **15**, 330. doi:10.1186/1471-2164-15-330
- Roman, J. (2006). Diluting the founder effect: cryptic invasions expand a marine invader's range. *Proc. R. Soc. B Biol. Sci.* **273**, 2453–2459. doi:10.1098/rspb.2006.3597
- Roman, J. and Palumbi, S. R. (2004). A global invader at home: population structure of the green crab, *Carcinus maenas*, in Europe. *Mol. Ecol.* **13**, 2891–2898. doi:10.1111/j.1365-294X.2004.02255.x

- Rossong, M. A., Quijón, P. A., Snelgrove, P. V. R., Barrett, T. J., McKenzie, C. H. and Locke, A.** (2012). Regional differences in foraging behaviour of invasive green crab (*Carcinus maenas*) populations in Atlantic Canada. *Biol. Invasions* **14**, 659-669. doi:10.1007/s10530-011-0107-7
- Sherman, C. D. H., Lotterhos, K. E., Richardson, M. F., Tepolt, C. K., Rollins, L. A., Palumbi, S. R. and Miller, A. D.** (2016). What are we missing about marine invasions? Filling in the gaps with evolutionary genomics. *Mar. Biol.* **163**, 198. doi:10.1007/s00227-016-2961-4
- Somero, G. N.** (2002). Thermal physiology and vertical zonation of intertidal animals: optima, limits, and costs of living. *Integr. Comp. Biol.* **42**, 780-789. doi:10.1093/icb/42.4.780
- Tepolt, C. K.** (2014). Adaptation and acclimation to temperature in a globally invasive marine species, *Carcinus maenas*. Doctoral Dissertation, Stanford University.
- Tepolt, C. K. and Palumbi, S. R.** (2015). Transcriptome sequencing reveals both neutral and adaptive genome dynamics in a marine invader. *Mol. Ecol.* **24**, 4145-4158. doi:10.1111/mec.13294
- Tepolt, C. K. and Somero, G. N.** (2014). Master of all trades: thermal acclimation and adaptation of cardiac function in a broadly distributed marine invasive species, the European green crab, *Carcinus maenas*. *J. Exp. Biol.* **217**, 1129-1138. doi:10.1242/jeb.093849
- Voss, E. R.** (2016). Conflicting geography in mitochondrial and nuclear markers in a green crab hybrid zone in the Gulf of Maine. *Honors Thesis*, Bowdoin College.
- Wade, M. J. and Brandvain, Y.** (2009). Reversing mother's curse: selection on male mitochondrial fitness effects. *Evolution* **63**, 1084-1089. doi:10.1111/j.1558-5646.2009.00614.x
- Willet, C. S.** (2010). Potential fitness trade-offs for thermal tolerance in the intertidal copepod *Tigriopus californicus*. *Evolution* **64**, 2521-2534. doi:10.1111/j.1558-5646.2010.01008.x
- Wolff, J. N., Ladoukakis, E. D., Enríquez, J. A. and Dowling, D. K.** (2014). Mitonuclear interactions: evolutionary consequences over multiple biological scales. *Proc. R. Soc. B. Biol. Sci.* **369**, 2013-0443. doi:10.1098/rstb.2013.0443