

Bowdoin College

Bowdoin Digital Commons

Biology Faculty Publications

Faculty Scholarship and Creative Work

9-1-2007

Disruption of topoisomerase II perturbs pairing in *Drosophila* cell culture

Benjamin R. Williams
Harvard Medical School

Jack R. Bateman
Harvard Medical School

Natasha D. Novikov
Harvard Medical School

C. Ting Wu
Harvard Medical School

Follow this and additional works at: <https://digitalcommons.bowdoin.edu/biology-faculty-publications>

Recommended Citation

Williams, Benjamin R.; Bateman, Jack R.; Novikov, Natasha D.; and Wu, C. Ting, "Disruption of topoisomerase II perturbs pairing in *Drosophila* cell culture" (2007). *Biology Faculty Publications*. 24. <https://digitalcommons.bowdoin.edu/biology-faculty-publications/24>

This Article is brought to you for free and open access by the Faculty Scholarship and Creative Work at Bowdoin Digital Commons. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of Bowdoin Digital Commons. For more information, please contact mdoyle@bowdoin.edu, a.sauer@bowdoin.edu.

Disruption of Topoisomerase II Perturbs Pairing in *Drosophila* Cell Culture

Benjamin R. Williams,* Jack R. Bateman,* Natasha D. Novikov* and C.-Ting Wu*^{†,1}

*Department of Genetics and [†]Division of Genetics and Division of Molecular Medicine, Harvard Medical School, Boston, Massachusetts 02115

Manuscript received May 22, 2007
Accepted for publication June 22, 2007

ABSTRACT

Homolog pairing refers to the alignment and physical apposition of homologous chromosomal segments. Although commonly observed during meiosis, homolog pairing also occurs in nonmeiotic cells of several organisms, including humans and *Drosophila*. The mechanism underlying nonmeiotic pairing, however, remains largely unknown. Here, we explore the use of established *Drosophila* cell lines for the analysis of pairing in somatic cells. Using fluorescent *in situ* hybridization (FISH), we assayed pairing at nine regions scattered throughout the genome of Kc₁₆₇ cells, observing high levels of homolog pairing at all six euchromatic regions assayed and variably lower levels in regions in or near centromeric heterochromatin. We have also observed extensive pairing in six additional cell lines representing different tissues of origin, different ploidies, and two different species, demonstrating homolog pairing in cell culture to be impervious to cell type or culture history. Furthermore, by sorting Kc₁₆₇ cells into G₁, S, and G₂ subpopulations, we show that even progression through these stages of the cell cycle does not significantly change pairing levels. Finally, our data indicate that disrupting *Drosophila topoisomerase II (Top2)* gene function with RNAi and chemical inhibitors perturbs homolog pairing, suggesting *Top2* to be a gene important for pairing.

ONE way in which the genome can be organized is through the physical pairing of homologous chromosomes. Such pairing occurs during meiosis, when it aligns chromosomes in preparation for recombination and segregation, as well as in nonmeiotic nuclei. Although most extensively studied in the Dipteran insect *Drosophila melanogaster*, nonmeiotic association of homologous chromosomal regions has also been observed elsewhere, where it participates in gene expression and other vital processes (reviewed by WU and MORRIS 1999; DUNCAN 2002; GRANT-DOWNTON and DICKINSON 2004; MCKEE 2004; ZICKLER 2006). Importantly, nonmeiotic pairing can influence gene regulation and DNA repair through the processes of transvection (reviewed by PIRROTTA 1999; WU and MORRIS 1999; DUNCAN 2002; KASSIS 2002; KENNISON and SOUTHWORTH 2002) and recombination (reviewed by GLOOR 2002; WYMAN *et al.* 2004; also see RONG and GOLIC 2003), respectively. Pairing of homologous chromosomal regions has also been implicated in mammalian X-inactivation (MARAHERNS 1999; BACHER *et al.* 2006; DIAZ-PEREZ *et al.* 2006; XU *et al.* 2006) and imprinting (LASALLE and LALANDE 1996; RIESSELMANN and HAAF 1999). Intriguingly, a reduced level of pairing at imprinted regions may be associated with syndromes such as autism and the Prader-Willi and Angelman syndromes (LASALLE and LALANDE 1996; THATCHER *et al.* 2005). Despite its role in multiple

phenomena, however, the mechanism(s) of nonmeiotic homolog pairing remains largely unknown.

To better understand nonmeiotic pairing, we have focused on *Drosophila*, where homologous chromosomes are essentially paired in all somatic cells throughout development (STEVENS 1907, 1908; METZ 1916; reviewed by DUNCAN 2002; MCKEE 2004). FISH analyses of embryos and tissues such as from the larval brain or imaginal discs indicate that, typically, a given chromosome region is paired in 60–100% of nuclei, but that this level of pairing can vary depending on the time of development, region being assayed, and phase of the cell cycle (KOPCZYNSKI and MUSKAVITCH 1992; CSINK and HENIKOFF 1998; FUNG *et al.* 1998; GEMKOW *et al.* 1998; SASS and HENIKOFF 1999; VAZQUEZ *et al.* 2002; RONSHAUGEN and LEVINE 2004; FRITSCH *et al.* 2006). For example, analysis of multiple chromosomal locations during early embryogenesis indicates that some regions initiate pairing earlier than do others, but that all regions eventually achieve high levels of pairing (FUNG *et al.* 1998). Other studies have examined the impact of the cell cycle on pairing. In one case, BrdU labeling of the larval central nervous system (CNS) revealed that the percentage of nuclei in which a euchromatic region, 59E, and a pericentromeric repeat, AACAC, were paired decreased during S phase from >75% to ~60% and from ~85% to <70%, respectively, and remained relatively low until after mitosis (CSINK and HENIKOFF 1998). In contrast, pairing at the internally repeated histone locus during the 13th embryonic mitotic cycle was found to be unperturbed by progression

¹Corresponding author: Department of Genetics, Harvard Medical School, 77 Ave. Louis Pasteur, NRB 264, Boston, MA 02115.
E-mail: twu@genetics.med.harvard.edu

through S phase and well into metaphase, decreasing only during anaphase, when the percentage of nuclei in which the locus was paired dropped from 63% to 19% (FUNG *et al.* 1998).

To date, efforts to identify genes that participate in somatic pairing have stemmed largely from observations of *in vivo* phenotypes that had been previously hypothesized to be sensitive to homolog pairing. For example, *zeste*, which encodes a transcription factor, was identified as a gene that modifies transvection-associated phenotypes of several loci (reviewed by PIRROTTA 1999; WU and MORRIS 1999; DUNCAN 2002; KENNISON and SOUTHWORTH 2002). Interestingly, *zeste* protein shows a propensity to self-aggregate, suggesting that it may promote pairing by binding to chromosomes and pulling or holding them together (CHEN and PIRROTTA 1993). However, it has yet to be determined how directly *zeste* affects pairing.

A more immediate connection has been made with *Polycomb* group (PcG) genes, which can mediate the pairing of PcG response elements (PREs) to effect long-range interactions and pairing-sensitive silencing (reviewed by PIRROTTA 1999; KASSIS 2002; KAVI *et al.* 2006; also see VAZQUEZ *et al.* 2006). For example, pairing of the PRE-containing element *Fab-7* requires several PcG proteins, such as Polycomb, Polyhomeotic, Posterior sex combs, and Polycomb-like (BANTIGNIES *et al.* 2003). A more recent study has shown that the pairing-sensitive silencing associated with *Fab-7* may also require members of the RNAi machinery, such as Dicer-2, PIWI, and Argonaute1 (GRIMAUD *et al.* 2006).

Another genetic element implicated in long-range interactions between homologous chromosomal regions is an insulator carried by the *gypsy* retrotransposon (reviewed by KUHN and GEYER 2003; CAPELSON and CORCES 2004; BRASSET and VAURY 2005; GASZNER and FELSENFELD 2006; VALENZUELA and KAMAKAKA 2006; also see KRAVCHENKO *et al.* 2005). Of note, two factors that work in conjunction with the *gypsy* insulator, Suppressor of Hairy Wing [Su(Hw)] and Modifier of *mdg4* [Mod(*mdg4*)] (KRAVCHENKO *et al.* 2005), colocalize at hundreds of chromosomal sites that do not correspond to *gypsy* elements and, along with another protein, Centrosomal Protein 190 (CP190), mediate the clustering of these sites into “insulator bodies” (reviewed by KUHN and GEYER 2003; CAPELSON and CORCES 2004; BRASSET and VAURY 2005; GASZNER and FELSENFELD 2006; VALENZUELA and KAMAKAKA 2006). Consistent with this finding, loss of Su(Hw) function has been observed to compromise homolog pairing (FRITSCH *et al.* 2006). Intriguingly, a protein that is associated with insulator function in mammals, the CCCTC binding factor (CTCF), has also been implicated in long-range interactions (LING *et al.* 2006; DONOHOE *et al.* 2007).

Despite these advances, we are still far from a full understanding of how homologs sense, find, and align with each other. To further our knowledge of somatic pairing

in *Drosophila*, we have extended our studies to cell culture in anticipation that such an experimental platform will facilitate analyses requiring the manipulation of homogenous populations of cells. A cell line-based approach could also allow researchers to identify genes involved in pairing solely on the basis of their capacity to support pairing, as assayed directly by visual examination. Importantly, because cell culture is unencumbered by developmental programs, it would permit the identification of genes that may also be essential for organismal viability.

Our study was encouraged by two reports. One, by CHERBAS and CHERBAS (1997), documented the capacity of a *Drosophila* immortal cell line to support paralogous recombination, a form of transformation attributed to somatic homolog pairing in which exogenous sequences integrate in the vicinity of homologous chromosomal regions. The second report, by HALFER and BARIGOZZI (1973), described two newly established *Drosophila* cell lines in which condensed mitotic chromosomes appeared entangled or aligned with their homologs in orcein- or quinacrine-stained spreads. These images were obtained with diploid as well as polyploid cell lines and recall earlier studies of Dipteran insects demonstrating the capacity of somatic pairing to accommodate polyploidy *in vivo* (METZ 1916, 1922, 1925; HOLT 1917). Interestingly, aligned condensed chromosomes have also been observed in cell lines of *Aedes* (mosquito), another Dipteran insect (NICHOLS *et al.* 1971). In our analysis, we used fluorescent *in situ* hybridization (FISH) to document somatic pairing of decondensed nonmitotic chromosomes in cell lines established decades ago and then characterized the consistency and breadth of that pairing. In addition, we obtained evidence indicating that *topoisomerase II* is a gene important for homolog pairing.

MATERIALS AND METHODS

Cell culture: Kc₁₆₇ (ECHALIER and OHANESSIAN 1969; PERRIMON 2007), S2 (SCHNEIDER 1972; PERRIMON 2007), D (DEBEC 1978, 1984; CHERBAS 2007), DH-33 (SONDERMEIJER *et al.* 1980; CHERBAS 2007), *mbn2* (HAARS *et al.* 1980; CHERBAS 2007), clone 8 (PEEL *et al.* 1990; PERRIMON 2007), and ML-DmBG2-c6 (U1 *et al.* 1994; CHERBAS 2007) cells were grown at 25° following standard protocols. Kc₁₆₇ and S2 cultures were grown in sterile filtered Schneider's medium [GIBCO (Grand Island, NY) no. 11720-034] supplemented with heat-inactivated fetal bovine serum (FBS, to a final concentration of 10% v/v; JRH 12103-78P) and penicillin–streptomycin (50 units/ml penicillin, 50 µg/ml streptomycin; GIBCO no. 15070-63). D, *mbn2*, DH-33, and ML-DmBG2-c6 cultures were grown in sterile filtered Shields and Sang M3 insect medium (S8523; Sigma, St. Louis) supplemented with FBS (10% v/v), BPYE [2.5 g/liter bactopectone (211677; Difco, Detroit) and 1.0 g/liter yeast extract (Y-1000; Sigma)], and penicillin–streptomycin (50 units/ml penicillin, 50 µg/ml streptomycin), with ML-DmBG2-c6 cultures being further supplemented with insulin (10 µg/ml; I6634, Sigma). Clone 8 cultures were grown in sterile filtered Shields and Sang M3 insect medium

supplemented with FBS (2% v/v), insulin (0.0125 IU/ml), fly extract (2.5% v/v; PERRIMON 2007), and penicillin–streptomycin (50 units/ml penicillin, 50 µg/ml streptomycin). To ensure that experiments were done with log-phase cells, active cultures were split at a 1:≥3 ratio, cultured for 1–2 days, and then passaged at $2\text{--}4 \times 10^6$ cells/ml prior to the analyses.

The Kc₁₆₇ and S2 cell lines were obtained from Norbert Perrimon; the D, DH-33, mbn2, and ML-DmBG2-c6 cell lines were obtained from the *Drosophila* Genome Resource Center; and the clone 8 cell line was obtained from Mitzi Kuroda.

FISH: Our FISH protocol was adapted from previously published protocols (MARSHALL *et al.* 1996; DERNBURG and SEDAT 1998; DERNBURG 2000; BANTIGNIES *et al.* 2003, 2005) and involved the following steps: Cells from log-phase cultures were adhered to either gelatin-coated (0.2%; G1393, Sigma) or lysine-treated (P8920; Sigma) 10-well glass slides (ER208W; Erie Scientific) for 1–3 hr. Slides were then gently washed with PBS (pH 7.2), fixed for 5 min with 4% formaldehyde in PBS (15700; Electron Sciences) at room temperature (RT), covered with a coverslip, frozen on an aluminum block (which had been precooled on dry ice), freed of their coverslips, and stored in 95% ethanol at -20° . After at least 20 min, slides were washed in $2\times$ SSCT (0.3 M NaCl, 0.03 M sodium citrate, 0.1% Tween-20)/formamide (5 min each in 0, 20, 40, and 50% formamide at RT and 30 min in 50% formamide/ $2\times$ SSCT at 37°). DNA probe in hybridization buffer was then added to the slides, covered with a coverslip, and denatured in an MJ Research (Watertown, MA) PTC-200 thermocycler with an Alpha Unit lock assembly block for 2 min at 91° , after which slides were transferred to a humidifying chamber, incubated overnight at $37^\circ\text{--}40^\circ$, and freed of their coverslips while being washed (30 min in 50% formamide/ $2\times$ SSCT at 37° , 5 min in 25% $2\times$ SSCT/formamide at RT, and three times in 5 min in $2\times$ SSCT at RT).

To visualize probes, slides were blocked in blocking buffer (0.1% BSA in $2\times$ SSCT) at RT for 30 min, incubated with either rhodamine-conjugated anti-DIG antibody (1207750; Roche, Indianapolis) or fluorescein anti-biotin (SP-3040; Vector, Burlingame, CA) in blocking buffer for 1.5 hr, and washed for 1 hr in $2\times$ SSCT, after which Vectashield with DAPI (H-1200; Vector) was added. Coverslips were applied and sealed to the slides with nail polish.

DNA probes were synthesized according to standard protocols. P1 plasmids (Berkeley *Drosophila* Genome Project) containing cloned *Drosophila* genomic DNA corresponding to chromosomal regions 21E3–4 (abbreviated as 21E3; DS03071; FUNG *et al.* 1998), 28B1–28B2 (abbreviated as 28B1; DS01529; FUNG *et al.* 1998), 40A2–40A3 (abbreviated as 40A2; DS09165; FUNG *et al.* 1998), and 69C2–69C8 (abbreviated as 69C2; DS02752; DEJ and SPRADLING 1999) were digested with restriction enzymes (DERNBURG 2000) and then labeled with either Digoxigenin (DIG-Nick translation mix, 1 745 816; Roche Diagnostics) or, for dual label experiments, Biotin (BioNick Labeling System, LT18247-015; Invitrogen) following the manufacturers' protocols. Probe for 16E1–16E2 (abbreviated as 16E1) was synthesized from the bacterial artificial chromosome BACR17D02 RP98-17D2 (AC012163; AE003507) by Nick translation/direct labeling (32-801300; Vysis) following the manufacturer's protocol. The 359-bp repeat probe was synthesized by PCR (DERNBURG 2000). Probes for 8C8 and 44F1 were synthesized from PCR products according to F. BANTIGNIES (personal communication): 8–10 1- to 1.4-kbp PCR products corresponding to genomic regions separated by ~ 1 kbp and spanning ~ 30 kbp were combined, purified, and labeled by Nick translation (FISH-Tag DNA kit; Invitrogen). Probes were diluted into hybridization buffer (50% formamide/ $2\times$ SSCT, 10% dextran sulfate) to a final concentration of $\sim 150\text{--}500$ ng/ $30\ \mu\text{l}$.

Oligo probes for the AACAC and dodeca heterochromatic repeats (DERNBURG and SEDAT 1998; DERNBURG 2000) were synthesized with either a 5' cy3 or cy5 fluorescent dye (Phoenix BioTechnologies) and contained locked nucleic acid (LNA) (SILAHTAROGLU *et al.* 2003, 2004) bases to increase melting temperature (AaCaCaAcAcAaCaCaAcAc and AcGgGgCaCgGg, for "AACAC" and "dodeca" probes, respectively, where uppercase letters denote LNA-modified nucleotides). An abbreviated FISH protocol was developed for LNA containing oligos: After fixation, cells were incubated for 30 min in $2\times$ SSCT at 37° , after which probe (1–100 nm in hybridization buffer) was added and slides were denatured at 91° for 2 min, washed immediately in $2\times$ SSCT for 30 min at 37° , and then mounted with Vectashield.

Microscopy: Through the generosity of Monica Colaiácovo, we were able to obtain images using a DeltaVision system with an Olympus IX-70 microscope and record data with a cooled CCD camera (model CH350; Roper Scientific). A U-APO 40 \times lens (NA 0.65–1.35) was used for all images except for those in Figure 1, where a PlanApo 60 \times lens (NA 1.4) was also used. Optical sections were collected at 0.50-µm increments and deconvolved using a conservative algorithm with 15 iterations. Data were analyzed with SoftWoRx Explorer software (Versions 1.0.1 and 1.1; Applied Precision). Three-dimensional images of nuclei were flattened for the figures.

Mitotic spreads: Chromosomes were spread using standard protocols (VEILE 1990; ECHALIER 1997), with or without incubating cells with colchicine (C3915; Sigma) for 2–24 hr.

Determining mitotic index: Anti- α -tubulin (T5168; Sigma) and secondary antibodies (SC-2010; Santa Cruz Biotechnology) were used following the manufacturers' instructions for labeling tissue culture cells. Mitotic cells were then identified by morphology. We confirmed the mitotic indexes corresponding to Figure 6, F and G (inset, Figure 6F) through assays using anti-phosphorylated-histone H3 (anti-PH3; 06-570; Upstate Biotechnology, Lake Placid, NY): control, 3.5% ($N = 2152$); camptothecin, 0.3% ($N = 1000$); *m*-amsacrine, 0.3% ($N = 1280$); and ICRF-193, 3.5% ($N = 976$).

Cell sorting: Live cells incubated with Hoechst 33342 (H3570; Molecular Probes, Eugene, OR) were sorted on a Vantage SE high-speed sorter (FACS Sorting Core Facilities; Children's Hospital, Boston), adhered to slides for 30–60 min, and then subjected to FISH. We confirmed that cells had been successfully sorted into G₁, S, and G₂ subpopulations by assessing nuclear DNA content as determined by DAPI staining (data not shown).

RNAi: Synthesis of dsRNA and application of RNAi to cells was carried out according to published protocols (CLEMENS *et al.* 2000; PERRIMON 2007). Control cells were treated with a blank of deionized water. Cells were fixed 3–4 days after treatment. Three distinct nonoverlapping dsRNAs were tested independently in studies of *Top1* as well as of *Top2*; outcomes did not vary among dsRNAs targeting the same gene.

Application of topoisomerase inhibitors: Our studies using camptothecin (C9911; Sigma; SOLIER *et al.* 2004), *m*-amsacrine (A9809; Sigma; SOLIER *et al.* 2004), and ICRF-193 (GR-332; Biomol; HOSSAIN *et al.* 2004) were guided by a sampling of studies carried out in mammals and *Drosophila*. Camptothecin and ICRF-193 were dissolved in DMSO (10 mM each), and *m*-amsacrine was dissolved in 30% ethanol (10 mM) before being added to log-phase cells to a final concentration of 10, 4, and 5 µM, respectively. Under the culture conditions used for this study, the doubling times for Kc₁₆₇ and clone 8 cells were estimated to be ~ 18 and ~ 15 hr, respectively. Accordingly, Kc₁₆₇ and clone 8 cells were incubated for 30 and 24 hr, respectively, except in the case of the clone 8 cells described in Figure 6, F and G, where, in one of the two trials, cells were incubated for an extended time of 30 hr. Control cells were treated with DMSO only.

Data analysis, percentile plots: All analyses were carried out two or more times, except for that of G₁ cells (Figure 5), which was conducted only once. To simplify the distance analyses for percentile plots, we assumed that cell lines with complex ploidies had the ploidy that was the most observed; for example, we assumed tetraploidy for Kc₁₆₇ cells, 94% of which had four copies of the X chromosome, four copies of chromosome II, and three to five copies of chromosome III. In support of this strategy, the conclusions of our analyses were not altered when the data were reanalyzed assuming other possible combinations of the observed ploidies. For nuclei with more than one signal, we identified signals likely to represent more than one copy of the targeted region by considering the expected ploidy for the nucleus, the number of signals obtained, and relative signal intensities. Often signal intensities did not permit us to assign copy number and, in these cases, we assigned copy number at random. We rarely observed Kc₁₆₇ nuclei (<1 of 2 × 10³) with more than four signals; instances when this did occur may have been due to background or represent G₂ nuclei with separated sister chromatids.

Nuclei that could not be scored due to lack of FISH signal, a weak signal-to-noise ratio, or high background were not included in the *N* values. With respect to assays that did not involve RNAi or chemical agents, the percentage of such nuclei varied from experiment to experiment. For example, Kc₁₆₇ and clone 8 cells yielded, respectively, 15 ± 8% (four trials) and 35 ± 15% (three trials) of nuclei that were not labeled in assays targeting 8C8, 15 ± 7% (two trials) and 9 ± 1% (two trials) in assays targeting 16E1, 7 ± 5% (seven trials) and 7 ± 10% (eight trials) in assays targeting 28B1, and 15 ± 11% (two trials) and 25 ± 2% (two trials) in assays targeting 44F1. Note that there was no obvious correlation between the degree of pairing and percentage of nuclei lacking signal in these cells (our unpublished observation).

The total percentage of nuclei without signal in Kc₁₆₇ cells treated with ICRF-193 was increased by an additional 2–27% (for example, a 27% increase would be one from 10 to 37%; three trials) in assays targeting 8C8, 12–14% (two trials) in assays targeting 16E1, 12–34% (five trials) in assays targeting 28B1, and 10–15% (two trials) in assays targeting 44F1. With regard to clone 8 cells treated with ICRF-193, we observed increases of 4–9% (three trials) in assays targeting 8C8, 12–16% (two trials) in assays targeting 16E1, 7–24% (six trials) in assays targeting 28B1, and 0–3% (two trials) in assays targeting 44F1. These observations also confirm the capacity of *Drosophila* cells to progress through mitosis in the presence of ICRF-193, consistent with other studies in which Top2 activity has been disrupted (CHANG *et al.* 2003; COELHO *et al.* 2003). Note that nuclei that are abnormally monosomic for a region targeted by FISH would artificially inflate the percentage of nuclei with single signals representing paired regions, suggesting that our assessment of the impact of ICRF-193 on pairing may be an underestimate.

To simulate a random distribution of FISH signals in the nucleus, we implemented a simple algorithm in Perl in which the nuclear shape of a typical Kc₁₆₇ nucleus was approximated as an ellipsoid defined by three radii (2.0, 2.0, and 2.5 μm; our unpublished observations), and the *x*, *y*, and *z* coordinates of any point within or on the surface of that ellipsoid were described by the following equation:

$$1 \geq \frac{x^2}{2.0^2} + \frac{y^2}{2.0^2} + \frac{z^2}{2.5^2}.$$

In our studies, we generated a random set of points (using the pseudorandom rand Perl function) with *x*, *y*, and *z* coordinates that fell in the ranges of [−2.0, 2.0] μm, [−2.0, 2.0] μm, and [−2.5, 2.5] μm, respectively. To model a tetraploid nucleus with

randomly placed FISH signals, we generated four points per ellipsoid and calculated the distances between them.

Matlab 7.0.1SP1 and Microsoft Excel 11.2 were used to analyze and plot the data. A two-tailed chi-square test was used to compare matched populations of nuclei with respect to the frequency of nuclei with a single signal. When experiments with more than one trial were compared, trials were combined. A two-tailed two-sample Kolmogorov–Smirnov test was used to compare the distances between populations displayed in percentile plots.

RESULTS

***Drosophila* immortal cell lines support pairing:** We first asked whether *Drosophila* immortal cell lines support homolog pairing by examining the commonly used *D. melanogaster* Kc₁₆₇ cell line, our subline of which is primarily tetraploid; among >50 nuclei (more than four trials), 94% had four copies of the X chromosome, four copies of chromosome II, and three to five copies of chromosome III (Table 1; Figure 1, A and B). To determine whether Kc₁₆₇ cells pair homologous chromosomes, we fixed log-phase cells with formaldehyde and then labeled them with fluorescent probes and used wide-field deconvolution microscopy to determine the positions of the signals in three-dimensional space. Regions targeted by the probes were considered paired when the center-to-center distance between FISH signals was ≤0.50 μm, the approximate distance at which adjacent FISH signals no longer overlap.

The first probe we used targeted region 28B1, located in a euchromatic portion of chromosome II (Figure 2A). Strikingly, 78.4 ± 8.4% of labeled nuclei produced a single FISH signal (*N* = 1507, nine trials; Figure 2, B and C). This high percentage of single-signal nuclei suggests that the majority of Kc₁₆₇ nuclei pair all four copies of the 28B1 region. To quantitate the range of distances between any two copies of 28B1, we assumed all nuclei to be tetraploid, calculated the distances between each pairwise combination of 28B1, as represented by FISH signals (Figure 2D, inset), and then arranged the distance data in a percentile plot (Figure 2D). Although distances between copies of 28B1 ranged from 0 μm to the full diameter of a nucleus (typically ~4–5 μm), 88% of all possible pairwise combinations were paired. Consistent with our observations of 28B1, significant pairing was also found for five other euchromatic regions (Figure 2, A, E, and F). We found that 72, 76, 82, 86, and 75% of nuclei gave a single signal when FISH was targeted, respectively, to 8C8 and 16E1 in the middle of the X, 21E3 and 44F1 at the end of the left arm and in the middle of the right arm of chromosome II, and 69C2 in the middle of the left arm of chromosome III. This consistent and high level of pairing across all six regions assayed is remarkable; regardless of probe used, between 72 and 86% of nuclei contained only a single signal even though there is variability in the efficacy of

TABLE 1
Pairing is observed in seven different cell lines

Line	Origin ^a	Year established	Karyotype ^b		% paired ^c
			II	III	
Kc ₁₆₇	Embryos	1969	4	3–5	78
S2	Embryos	1972	4, 6	4, 6	76
D	Embryonic haploid cells	1978	2, 4	2, 4	83
DH-33	Embryos, <i>D. hydei</i>	1980	4	4–5	82
mbn2	Hemocytes	1980	ND ^d	ND ^d	69
Clone 8	Imaginal wing discs	1990	2	2	87
ML-DmBG2-c6	Larval central nervous system	1994	3 ^e	2 ^e	86

See Figure 1 for examples of mitotic figures and MATERIALS AND METHODS for references for cell lines.

^aAll cells lines were from *D. melanogaster*, except as noted for DH-33.

^b $N > 50$ for Kc₁₆₇, $N = 20$ for clone 8, and $N > 7$ for all other lines except mbn2 (ND, not determined).

^cData represent the percentage of single-signal nuclei as assayed at the 28B1 region (Figure 4A except in the case Kc₁₆₇, where the data correspond to Figure 2C); $N > 55$ for each line.

^dAssumed tetraploid.

^eAberrant chromosome morphology (Figure 1, G and H).

FISH, as determined by signal strength, background staining, and the percentage of nuclei lacking signal (MATERIALS AND METHODS).

Using the percentile plot, we then assessed the statistical significance of the pairing we observed. First, we determined the frequency at which two or more of the four copies of 28B1 would be expected to colocalize solely by chance by computationally simulating three-dimensional nuclei containing four randomly placed FISH signals. Virtually no computer-simulated nuclei colocalized all four signals. In fact, 90% of these nuclei showed four completely separated signals. In addition,

percentile plots revealed that only 2% of all pairwise combinations of signals were colocalized (black line, Figure 3A). In our second approach, we applied dual-labeled FISH to experimentally establish the frequency of colocalization for two regions that do not share homology. Here, we measured the distances between two sets of two loci each: 21E3 and 28B1, both on chromosome II, and 28B1 and 69C2, on chromosomes II and III, respectively (Figure 3B). Colocalization was observed for only 5 and 0.8% of all pairwise combinations of the 21E3 and 28B1 regions and the 28B1 and 69C2 regions, respectively (Figure 3B). These data demonstrate that




FIGURE 1.—Karyotypes and chromosomal pairing (40–96 \times magnification). (A) Kc₁₆₇ mitotic figure stained with DAPI. Chromosome IV is not noted. (B) Kc₁₆₇ mitotic figures labeled with DAPI (blue) and probes targeting the AACAC (green, chromosome II) and dodeca (red, chromosome III) pericentric heterochromatin repeats confirm four copies of X and chromosome II and three to five copies of chromosome III. The X chromosome was identified by its unique morphology. (C–F) Indicative of pairing, chromosomes (red lines) of similar morphology are occasionally intertwined or grouped in Kc₁₆₇ (C and D) and clone 8 (E and F) mitotic figures. Clone 8 cells have a single X, a single Y, and two copies each of chromosomes II and III; chromosome IV is not noted. (G and H) ML-DmBG2-c6 mitotic figures stained with DAPI (G) or labeled with AACAC (yellow) and dodeca (red, H) include chromosomes with aberrant morphology.




FIGURE 2.—*Drosophila* cell culture supports pairing. (A) Regions targeted by probes. Chromosome IV is not shown. (B) Flattened images of Kc_{167} nuclei stained with DAPI and FISH targeting 28B1. (C) Percentage of Kc_{167} nuclei \pm SD (nine trials) with the indicated number of signals per nucleus identifying 28B1. (D) Percentile plot of data shown in C, assuming tetraploidy. All inter-signal distances within each nucleus are ranked, shortest to longest, left to right, where 100% corresponds to the greatest distance observed. Signals separated by $\leq 0.50 \mu\text{m}$ (---) are considered one signal. Open circles, data from inset illustrating distance calculations of nucleus with three signals. (E and F) Pairing profile and percentile plot when FISH was targeted to euchromatic regions. (G and H) Pairing profile and percentile plot when FISH was targeted to regions in or near heterochromatin. In this and all other figures, bars, $5 \mu\text{m}$; N, number of nuclei.

euchromatic regions rarely overlap by chance, strongly reinforcing our observation that somatic homolog pairing can occur genomewide and be maintained in transformed cell lines. Importantly, the amount of pairing we obtained approximates that observed *in vivo* (CSINK and

HENIKOFF 1998; FUNG *et al.* 1998; GEMKOW *et al.* 1998; SASS and HENIKOFF 1999). As Kc_{167} cells represent a departure from diploidy, these data further indicate that somatic pairing can tolerate levels of aneuploidy that are typically found in established cell lines.




FIGURE 3.—(A) Percentile plot for randomly placed FISH signals in theoretical nuclei with radii of 2, 2, and 2.5 μm compared to the percentile plot of an experiment in which Kc_{167} nuclei were labeled with probe targeting 28B1. Distances between randomly placed FISH signals are significantly greater than those between signals representing 28B1 ($P < 1 \times 10^{-10}$). (B) Percentile plot of distances between signals representing 21E3 and 28B1 and between signals representing 28B1 and 69C2. Data from A are added for comparison. Distances between signals representing 21E and 28B1 or 28B1 and 69C2 are significantly greater than those between signals representing 28B1 in A ($P_s < 1 \times 10^{-10}$).

In contrast to our findings at euchromatic regions, we found significantly less pairing at regions located in or near the centromeric heterochromatin of all three major chromosomes ($P \leq 0.05$; Figure 2, G and H). Using probes targeting a 359-bp repeated element of the centromeric heterochromatin of X, region 40A2 located just outside the centromeric heterochromatin of chromosome II, and the dodeca-repeated element located in the centromeric heterochromatin of chromosome III (Figure 2A), we observed only 24, 51, and 13%, respectively, of nuclei giving a single signal (Figure 2G) and correspondingly modest pairing levels of 45, 67, and 49% (Figure 2H). While it is possible that this lower level of pairing resulted not from the unpaired state of homologs but rather from the potential of repeated elements to form multiple distinct clusters, the apparent reduction in pairing we observed when visualizing unique sequences in the pericentromeric region of 40A2 argues against this as the sole interpretation. Interestingly, HALFER and BARIGOZZI (1973) also observed that the heterochromatic regions of condensed mitotic chromosomes were less paired than their euchromatic counterparts. Note that as mitotic cells represented only $\sim 2.1\%$ (see below) of our cultures, our findings cannot be explained solely by the behavior of mitotic chromosomes.

Homolog pairing occurs in a variety of cell lines: The widespread and persistent nature of homolog pairing in Kc_{167} cells encouraged us to ask whether the mechanism underlying pairing has also been maintained in other cell lines and, if so, whether the degree of pairing would

vary with cell type or culture history. Accordingly, we assayed pairing at 28B1 in six additional established cell lines (Table 1). These lines differ in terms of their species of origin (*D. melanogaster* and *D. hydei*), morphology, tissue of origin (embryonic, epithelial, hemocyte, imaginal disc, and CNS), year of establishment (1969–1994), and karyotype/ploidy (diploid, polyploid/aneuploid). They include the popular S2 cell line (SCHNEIDER 1972), which was derived from late-stage embryos and, like its derivative sublines such as S2R+ (YANAGAWA *et al.* 1998), is used in a wide variety of studies. The S2 cells we used proved to be generally tetraploid. In contrast, the male (*X/Y*) clone 8 cell line, which originated from imaginal wing discs, is diploid (Table 1) (PEEL *et al.* 1990; PEEL and MILNER 1991). The D cell line is also noteworthy, as it was established from haploid embryos (DEBEC 1978, 1984), even though it is currently a mixed population of diploid and tetraploid cells (Table 1); as such, there is no history of *in vivo* homolog pairing for chromosomes of the D cell line. Following standard protocols (MATERIALS AND METHODS), aliquots of each of the cell lines were cultured to generate log-phase cells. Using probes targeting 28B1, we found a nearly uniform high level of pairing that was reminiscent of that observed in Kc_{167} cells (Figure 4).

Homolog pairing remains equally strong in G_1 , S, and G_2 : Taking advantage of the amenability of cell culture to cell sorting, we addressed the impact of cell cycle on pairing. Log-phase Kc_{167} cells incubated in Hoechst dye were sorted into G_1 , S, and G_2 subpopulations on the basis of fluorescence intensity. Sorted cells




FIGURE 4.—Seven different cell lines support pairing. (A) Pairing profiles and (B) percentile plots of nuclei imaged with FISH directed against 28B1.




FIGURE 5.—Pairing is maintained through G₁, S, and G₂ phases of the cell cycle. (A) Pairing profiles of G₁, S, G₂, and unsorted Kc₁₆₇ cells imaged with FISH targeting 28B1. Analysis of G₂, S, and unsorted cells was repeated with similar results (not shown). (B) Percentile plot of data in A showing that distance distributions of G₁, S, G₂, and unsorted cells do not differ significantly from each other ($P_s \geq 0.95$). Inset, cells were stained with Hoechst 33342 and sorted by DNA content.

were then allowed to adhere to glass slides for ~30 min in medium prior to being fixed in formaldehyde and then subjected to FISH analysis. Our data indicate that the level of pairing, as assayed at 28B1, is uniform across all three subpopulations (Figure 5) and does not differ significantly from that of unsorted cells ($P_s \geq 0.95$); the slightly lower level of pairing observed for unsorted cells may reflect its inclusion of mitotic cells.

Although we cannot rule out that the observed pairing derives from a potential technical bias that enriches for cells in which homologs are paired, it is noteworthy that our observation is in line with the report by FUNG *et al.* (1998) that pairing can be maintained through S phase and into metaphase. A constant level of pairing in the G₁, S, and G₂ portions of the cell cycle differs, however, from the disruption of pairing described by CSINK and HENIKOFF (1998) in S-phase larval CNS cells. These diverse patterns of pairing may reflect different sensitivities of pairing to cell type, ploidy, and/or length of the cell cycle (GOLIC and GOLIC 1996; GUBB *et al.* 1997).

To examine mitotic chromosomes, we treated cells with a hypotonic solution and then fixed them in a mix of acetic acid and methanol (ECHALIER 1997). In this way, we were occasionally able to observe an intertwining of chromosomes in Kc₁₆₇ cells (Figure 1, C and D) as well as unambiguous alignments of homologs as fully condensed mitotic chromosomes, in the form of paired dyads, in clone 8 cells (Figure 1, E and F). These findings are reminiscent of the homolog entanglements and alignments characterized *in vivo* by STEVENS (1907, 1908), METZ (1916, 1922, 1925), and HOLT (1917) and in *Drosophila* cell lines by HALFER and BARIGOZZI (1973), as well as in *Aedes* cell lines by NICHOLS *et al.* (1971). They demonstrate that homologous chromosomes have a capacity to remain paired in at least some stages of mitosis and are consistent with FISH analyses showing that pairing is not perturbed at the histone locus during the 13th embryonic mitosis until after metaphase (FUNG *et al.* 1998). That M-phase pairing is tenacious is furthered by its ability to withstand hypotonic treatment.

Disruption of Top2 perturbs homolog pairing:

Taken together, our data establish that pairing in cell culture approximates pairing *in vivo* in terms of level and tenacity. This finding suggests that *Drosophila* cell

culture can serve as a powerful experimental system for identifying genes responsible for somatic homolog pairing. In particular, FISH-based screens in cell culture would enable direct visual assessments of pairing in different RNAi-driven mutant backgrounds. In fact, focusing our analysis on euchromatic regions, where homolog pairing is maximal, we have found that dsRNA targeting *Drosophila Top2* expression can reduce the level of pairing in Kc₁₆₇ cells.

Top2 encodes the single *Drosophila* type II DNA topoisomerase, which represents a class of enzymes that generate double-strand breaks (DSBs), can catenate and decatenate DNA, and are well known for their role in chromosome structure and condensation and sister chromatid segregation (reviewed by PORTER and FARR 2004; also see CHANG *et al.* 2003; COELHO *et al.* 2003; SAVVIDOU *et al.* 2005; SMILEY *et al.* 2007). We have found that treatment of log-phase Kc₁₆₇ cells with any one of three dsRNAs directed against Top2 function followed by FISH targeting 28B1 reduced the percentage of nuclei with a single signal from $87 \pm 1\%$ to $55 \pm 5\%$ ($P < 0.0001$; Figure 6, A and B) and lowered pairing levels from 91 to 71% ($P < 0.0001$; Figure 6C).

To extend these findings, we determined the impact of two chemical agents that block topoisomerase II: ICRF-193, which stabilizes a noncovalent complex between the enzyme and DNA without generating DSBs, and *m*-amsacrine, which leads to the accumulation of DSBs (reviewed by ANDOH and ISHIDA 1998; LARSEN *et al.* 2003). Treatment with ICRF-193 (10 μM) reduced the percentage of nuclei with a single signal from $84 \pm 4\%$ to $54 \pm 4\%$ ($P < 0.0001$; Figures 6D and 7; also see Table 2, which includes data not shown in Figure 6D). This finding is unlikely to reflect an increase in the percentage of cells in M phase as the mitotic index of ICRF-193-treated cells is, if anything, lower than that of untreated controls (2.1 and 1.0%, respectively; Figure 6D, inset; CHANG *et al.* 2003; COELHO *et al.* 2003). In contrast to treatment with ICRF-193, treatment with *m*-amsacrine (5 μM) had much less of an effect (Figures 6D and 7). This result is consistent with the fact that ICRF-193 and *m*-amsacrine have different mechanisms of action (reviewed by ANDOH and ISHIDA 1998; LARSEN *et al.* 2003), although we cannot rule out a technical basis for the ineffectiveness of *m*-amsacrine. Treatment




FIGURE 6.—Disruption of Top2 perturbs homolog pairing. (A–C) dsRNA directed against Top2 reduces pairing at 28B1 in *Kc*₁₆₇ cells as illustrated by nuclei imaged with DAPI and FISH (A), a pairing profile (B), and a percentile plot (C). dsRNA directed against Top1 has little effect. B shows the results (\pm SD) of experiments using three distinct dsRNAs targeting Top2 and two distinct dsRNAs targeting Top1, as well as two control experiments run without dsRNA. The percentile plot combines data described in B. (D and E) ICRF-193 (10 μ M) reduces pairing at 28B1 (D) and 16E1 (E) in *Kc*₁₆₇ cells. Camptothecin (4 μ M) and *m*-amsacrine (5 μ M) are less perturbing. Control nuclei were from cells treated with DMSO. Data represent two trials. (F and G) Pairing is more disrupted at 28B1 (F; $P < 0.0001$) than it is at 16E1 (G; $P = 0.003$) by ICRF-193 in clone 8 cells. Little disruption was observed when cells were treated with camptothecin (4 μ M), *m*-amsacrine (5 μ M), or DMSO (control). The data are from two independent experiments, one in which cells were treated with drugs for 24 hr and the other in which cells were treated for 30 hr. For each time point, nuclei imaged with FISH targeting 28B1 or 16E1 were from the same culture. *N*, number of nuclei from both time points combined. (Insets in D and F) Mitotic indexes correspond to one of the two independent trials shown.




FIGURE 7.—ICRF-193 disrupts pairing in Kc_{167} cells. Nuclei are from Kc_{167} cells treated with inhibitors of topoisomerase I ($4 \mu\text{M}$ camptothecin) or topoisomerase II ($5 \mu\text{M}$ *m*-amsacrine, $10 \mu\text{M}$ ICRF-193) and imaged with FISH targeting 28B1. Control nuclei are from cells treated with DMSO. Note the nuclei of cells treated with ICRF-193 can be misshapen. Nuclei are from cultures represented in Figure 6, D and E.

with dsRNA (Figure 6, A–C) or an enzymatic inhibitor, camptothecin ($4 \mu\text{M}$), directed against a different topoisomerase, topoisomerase I (Top1; reviewed by POMMIER 2006), had similarly minimal effects in our assays (Figures 6D and 7). Finally, we demonstrated that the impact of ICRF-193 was not specific for 28B1 by targeting FISH to 8C8, 16E1, and 44F1. We observed a decrease in the percentage of single-signal nuclei from $73 \pm 2\%$ to $55 \pm 2\%$, $65 \pm 7\%$ to $32 \pm 4\%$, and $79 \pm 10\%$ to $59 \pm 9\%$, respectively, suggesting that the effect of ICRF-193 on pairing may be general, although variable, across euchromatic regions (Table 2, $P_s < 0.0001$; Figure 6E).

These observations must be considered in light of the cell cycle and the two forms of homologous pairing that come into play: pairing between homologs (including the pairing of subchromosomal regions or fragments thereof) and pairing, or cohesion, between sister chromatids (Figure 8A). *A priori*, reduced pairing in G_1 prior to the synthesis of sister chromatids would necessarily imply the unpaired state of homologs. In contrast, reduced pairing in G_2 could result from the unpaired state of either homologs or sister chromatids or some combination thereof (see Figure 8 legend for further discussion). The data presented thus far do not distinguish between these possibilities.

Equally important for consideration is that mitosis in the presence of Top2 inhibitors can cause sister chromatids to cosegregate to the same daughter nucleus (not shown in Figure 8; reviewed by PORTER and FARR 2004) and produce cells that are monosomic, nullisomic, or segmentally aneuploid. Indeed, treatment with ICRF-193 generated a significant level of aneuploidy; we observed that the total percentage of nuclei lacking FISH signal can be increased by an additional 34% (for example, from 3 to 37%), indicating that the mitotic divisions producing these nuclei had cosegregated all copies of the targeted chromosomal region to only one daughter nucleus (see MATERIALS AND METHODS for additional information). As such, an increase in the percentage of nuclei with multiple FISH signals from cultures treated with ICRF-193 could also reflect the separation in G_1 aneuploid cells of what were once sister chromatids.

To decipher what form of unpairing underlies the effect of ICRF-193, we turned to male (X/Y) clone 8 cells where, *a priori*, lack of a homolog for the single X ($N = 20$ cells; Table 1) should allow the unpaired state of homologs to be more easily distinguished from sister chromatid separation (Figure 8B). Specifically, conditions that disrupt only homolog pairing should decrease the number of single-signal nuclei when FISH is targeted to autosomal regions but not when FISH is targeted to X-linked regions. In contrast, conditions that separate sister chromatids in G_2 and/or in aneuploid G_1 cells should decrease the number of single-signal nuclei regardless of whether FISH targets autosomal or X-linked regions.

We found that application of ICRF-193 to clone 8 cells reduced the percentage of single-signal nuclei by only 6.9% (of total nuclei scored) when FISH was targeted to 16E1 on the X even as it reduced the percentage by 37.8% when FISH was targeted to autosomal 28B1 ($P = 0.02$ and $P < 0.0001$, respectively; data included in Figure 6, F and G, and Table 2; Figure 9). Aiming to augment the impact of ICRF-193 at 16E1, we repeated our analysis while increasing the exposure time of the cells to ICRF-193. Again, we observed a much smaller effect when FISH was targeted to 16E1 as compared to 28B1, with the reduction of single-signal nuclei being 4.5 and 39.7%, respectively ($P = 0.08$ and $P < 0.0001$,

TABLE 2
ICRF-193 disrupts pairing more effectively in the presence of a homolog

	8C8 (%)		16E1 (%)		28B1 (%)		44F1 (%)	
	Clone 8	Kc ₁₆₇	Clone 8	Kc ₁₆₇	Clone 8	Kc ₁₆₇	Clone 8	Kc ₁₆₇
Control	<u>93 ± 2</u>	73 ± 2	<u>89 ± 1</u>	65 ± 7	84 ± 4	84 ± 5	90 ± 6	79 ± 10
ICRF-193	<u>86 ± 2</u>	55 ± 2	<u>83 ± 1</u>	32 ± 0.4	51 ± 5	52 ± 5	75 ± 3	59 ± 9
Difference	<u>7 ± 3</u>	18 ± 2	<u>6 ± 1</u>	33 ± 7	31 ± 6	32 ± 6	15 ± 7	20 ± 13

Data represent the percentage of single-signal nuclei (\pm SD). N s = 149–583 (two to three trials), except in the case of Kc₁₆₇ cells treated with ICRF-193 and labeled with probe targeting 16E1, where $N = 93$ (two trials). Underlining indicates data from FISH analyses targeting regions on the X chromosome in clone 8 cells, which carry only a single X.

respectively; data included in Figure 6, F and G, and Table 2). Similar results were obtained when we targeted FISH to another X-linked region, 8C8, and another autosomal locus, 44F1; here we observed that single-signal nuclei were reduced by $7 \pm 3\%$ and $15 \pm 7\%$,

respectively (P s < 0.0001; Table 2). In short, although we observed small reductions in the percentage of single-signal nuclei when FISH was targeted to regions on the X, we observed greater reductions when FISH was targeted to autosomal regions. Importantly, this lesser effect at 16E1 and 8C8 is not due to an insensitivity of these regions to ICRF-193, as pairing at 16E1 and 8C8 in Kc₁₆₇ cells is as susceptible to ICRF-193 as is pairing at autosomal regions (Figure 6, D and E; Table 2). These findings indicate that while sister chromatid separation may be occurring, it is unlikely to fully explain the effect of ICRF-193, arguing that loss of Top2 activity can disrupt homolog pairing.

As mentioned just above, our data suggest that loss of Top2 function may also disturb sister chromatid cohesion to a limited extent. Consistent with this, we have found that treatment with ICRF-193 augments the number of nuclei whose single signals actually consist of two fluorescent spots that are, however, too closely spaced ($\leq 0.50 \mu\text{m}$) to be considered separate (MATERIALS AND METHODS). The total percentage of nuclei with such closely spaced spots was increased by an additional 1–11%




FIGURE 8.—Use of clone 8 cells to distinguish unpaired homologs from separated sister chromatids. (A) In tetraploid Kc₁₆₇ cells, which carry four X chromosomes, multiple signals in a nucleus may represent unpaired homologs in either G₁ or G₂ or separated sister chromatids in G₂, regardless of whether they identify X-linked or autosomal regions. (B) In clone 8 cells, which carry only one X chromosome, multiple signals identifying X-linked regions can represent only separated sister chromatids in G₂. In contrast, multiple signals identifying autosomal regions can represent unpaired homologs in either G₁ or G₂ or separated sister chromatids in G₂. Note that the number of homologs or sister chromatids that are unpaired can differ from that shown. Variation of signal intensity did not allow accurate experimental determination of copy number per signal (data not shown). Also note that homolog pairing could occur between the sister chromatid of a chromosome and the homolog of that same chromosome. As such, the forms of sister chromatid separation illustrated may also entail an unpairing of homologs. If so, any separation of sister chromatids will necessarily imply homolog unpairing. This figure does not address situations involving aneuploidy. N/A, sister chromatids do not exist prior to replication.




FIGURE 9.—ICRF-193 disrupts pairing in clone 8 cells. Nuclei are from clone 8 cells treated with inhibitors of topoisomerase I (4 μ M camptothecin) or topoisomerase II (5 μ M *m*-amsacrine, 10 μ M ICRF-193) and imaged with FISH targeting 28B1 (A) or 16E1 (B). Control nuclei are from cells treated with DMSO. Nuclei are from cultures represented in Figure 6, F and G.

(three trials; for example, an 11% increase would be one from 1 to 12%) and 6–8% (two trials) in clone 8 cells when FISH was targeted to 8C8 and 16E1, respectively, and 0–11% (two trials) and 2–10% (five trials) when FISH was targeted to 28B1 and 44F1, respectively. This observation suggests that ICRF-193 may antagonize sister chromatid cohesion in G_2 nuclei in addition to disrupting homolog pairing.

DISCUSSION

In this study, we demonstrated that *Drosophila* immortal cell lines can support homolog pairing. Using FISH, we found remarkably consistent and high levels of pairing in tetraploid Kc_{167} cells and observed that such levels equally characterize G_1 , S, and G_2 cells. Strong pairing was also seen in six additional *Drosophila* cell lines representing two species and different tissue types and ploidies. Of particular note is the pairing that occurs in the D cell line. As this line originated from haploid cells, becoming a mix of diploid and tetraploid cells only after having become established (Table 1) (CHERBAS 2007), the pairing observed in D cells indicates that homolog pairing can initiate *ex vivo* and does not require homologs to be of different parental origin.

We also observed homolog pairing during M phase. Such homolog alignment is reminiscent of meiotic pairing and may reflect a commonality between the two forms of pairing. It further suggests that homolog interactions may be able to endure cell division, perhaps via the persistence of chromatin marks, physical associations, nuclear territories (FUNG *et al.* 1998; BANTIGNIES *et al.* 2003), or perichromosomal layers (VAN HOOSER *et al.* 2005).

Theoretically, enduring associations are compatible with mitosis. For instance, if each sister chromatid of a G_2 dyad is paired with one sister of the homologous dyad to produce two sets of paired homologs, pairing could be maintained through mitosis if paired homologs cosegregate. In fact, assuming sister chromatid separation of one dyad to be random with respect to that of its homologous dyad, paired homologs will as often cosegregate as they will be pulled apart (FUNG *et al.* 1998). This scenario predicts a loss of pairing during mitosis, but not a complete loss, and is consistent with observations of embryonic mitoses (FUNG *et al.* 1998). Note, however, that mitosis could also progress without fully unpairing any set of homologs while still accommodating the unpairing observed during mitosis. For example, if homologs were permanently attached but at only a few or just one locus, regions elsewhere on the chromosomes would be free to unpair. Alternatively, if paired homologs were permanently attached via one strand of DNA from each homolog such that replication produced two dyads that are paired via only one sister per dyad, mitoses would generate a mix of paired and unpaired homologs, the latter being expected to become paired in the subsequent G_1 . Mechanisms such as these raise the possibility of immortal chromosome attachments and recall observations of immortal DNA strands in asymmetrically dividing cells (reviewed by CAIRNS 2006; see also CONBOY *et al.* 2007). Finally, we note that mitosis may be a means by which pairing and pairing-sensitive phenomena are regulated; if there are instances when pairing must be maintained, mitotic mechanisms could orchestrate paired homologs at the metaphase plate such that they cosegregate while, if ever pairing must

be destroyed, the cell could trigger a mitosis in which paired homologs are forced apart.

Our studies also revealed that pairing of regions in or near centric heterochromatin is significantly less than that observed at euchromatic regions. This finding is consistent with observations of condensed mitotic chromosomes by HALFER and BARIGOZZI (1973) and could be explained if heterochromatic regions pair less, unpair more, or pair more slowly than do euchromatic regions. The prevalence of repeated sequences in heterochromatin may contribute to the unpaired state of homologs by, for example, promoting intrachromosomal pairing and/or homolog misalignment. Our data are also consistent with a loss of sister chromatid cohesion in centromeric regions. This explanation, however, would not be in line with studies showing that cohesin proteins, which hold sister chromatids together, persist at the centromeres longer than they do along chromosome arms (reviewed by WANG and DAI 2005; WATANABE 2005). As such, our data may indicate that the lower degree of homolog pairing observed in or near pericentric heterochromatin in *Kc*₁₆₇ cells results from a preoccupation of centromeric regions with sister chromatid, *vs.* homolog, pairing.

This report also summarizes our identification of *Top2* as a gene important for homolog pairing. Here, we found that loss of *Top2* function reduced pairing to a greater degree at autosomal *vs.* X-linked regions in diploid male clone 8 cells. This finding indicates that the reduction in pairing cannot be explained solely by a loss of cohesion between sister chromatids and argues that *Top2* plays a role in homolog pairing. How might *Top2* influence homolog pairing? It might act indirectly through its role in cell cycle events, such as chromosome condensation, congression of chromosomes at the metaphase plate, and separation of sister chromatids during mitosis (reviewed by PORTER and FARR 2004; also see CHANG *et al.* 2003; COELHO *et al.* 2003; HOSSAIN *et al.* 2004; SAVVIDOU *et al.* 2005). For example, cosegregation of sister chromatids due to a lack of *Top2* activity may generate a degree of disorganization that is incompatible with homolog pairing. Alternatively, abnormal presence of a sister chromatid may prohibit a chromosome from pairing with its homolog. That sister chromatid and homolog pairing may be mutually exclusive raises the possibility that the two are mutually regulated and may even explain how the genome can orchestrate sister chromatid separation during mitosis even as it promotes homolog pairing in the ensuing interphase. Interestingly, as sister chromatid cohesion has been implicated in gene regulation (AZUARA *et al.* 2003; MLYNARCZYK-EVANS *et al.* 2006; also see DORSETT 2007), an interplay between homolog pairing and sister chromatid cohesion could also provide a mechanism by which homolog pairing can modulate gene regulation.

Top2 may also have a direct role in the paired state of homologs. For example, it may bring and/or hold ho-

mologs together by altering DNA topology, modifying chromatin or chromosome structure, or catenating DNA molecules (BACHANT *et al.* 2002; VAZQUEZ *et al.* 2002; IWABATA *et al.* 2005). Alternatively, it may dimerize while bound to DNA or via its role as a chromosome structural/scaffold protein (EARNSHAW and HECK 1985; EARNSHAW *et al.* 1985; GASSER *et al.* 1986; more recently MAESHIMA and LAEMMLI 2003; MAESHIMA *et al.* 2005); for instance, it may act in conjunction with Su(Hw) (FRITSCH *et al.* 2006), whose DNA binding site contains the *in vitro* recognition sequence for topoisomerase II (NABIROCHKIN *et al.* 1998). Interestingly, if homolog pairing is required for the proper arrangement and segregation of chromosomes during mitosis, such unpairing could have contributed to the aneuploidy we observed in the presence of ICRF-193.

Top2 may also influence pairing via its capacity to decatenate chromosomes; just as the decatenation activity of *Top2* is required for the separation of sister chromatids, so may it be required prior to or during mitosis for the separation of paired homologs, should such pairing involve catenation. In this case, loss of *Top2* would lead to the cosegregation of catenated homologs, whose entanglement may then inhibit homolog pairing, and perhaps even sister chromatid cohesion, in the daughter nuclei.

Finally, it is worth considering the role of topoisomerase II activity in transcription (JU *et al.* 2006), where it appears to generate transient DSBs during gene activation. *Top2* may therefore affect pairing via its impact on chromatin structure during the transcriptional process. This interpretation is reminiscent of the implication of transcription, and the regulation thereof, in pairing (reviewed by MCKEE 2004) and the association of *Top2* with PcG target sequences and Polyhomeotic (LUPO *et al.* 2001), both of which participate in pairing-sensitive gene regulatory phenomena (reviewed by PIRROTTA 1999; KASSIS 2002; KAVI *et al.* 2006).

Our findings are consistent with proposals suggesting that topoisomerase II may contribute to pairing during meiosis (COBB *et al.* 1997; VAZQUEZ *et al.* 2002; IWABATA *et al.* 2005). In this regard, it is of interest that *Spo11*, which is homologous to a type II DNA topoisomerase, promotes meiotic pairing through the generation of DSBs in yeast, mice, and plants (reviewed by MCKEE 2004; GERTON and HAWLEY 2005; PAWLOWSKI and CANDE 2005; COLAIACOVO 2006; ZICKLER 2006). As such, the role of *Top2* in somatic pairing may be analogous to that of type II topoisomerases in meiotic pairing. In fact, a contribution of *Top2* to pairing in nonmeiotic cells may explain why *Drosophila* meiosis, which is believed to arise from the pairing that occurs in premeiotic cells (reviewed by MCKEE 2004; GERTON and HAWLEY 2005; ZICKLER 2006), does not require the generation of meiotic DSBs.

Curiously, absence of *Spo11*, as well as other genetic backgrounds, can lead to the meiotic pairing of

nonhomologous chromosomes (McCLINTOCK 1933; reviewed by MCKEE 2004; GERTON and HAWLEY 2005; PAWLOWSKI and CANDE 2005; COLAIACOVO 2006; OSMAN *et al.* 2006; ZICKLER 2006; also see TSUBOUCHI and ROEDER 2005), which was proposed by McCLINTOCK (1933) to be competitive with homologous pairing. Hence, loss of somatic pairing brought about by disrupting Top2 activity may be accompanied, maintained, or even driven by nonhomologous pairing. The concept that homologous and nonhomologous pairing are mutually exclusive suggests that interplay between them may provide a general mechanism for modulating nuclear organization and gene expression. Indeed, interactions between nonhomologous chromosomal regions may be as potent as those between homologous regions and, as may be the case with homologous pairing, have the capacity to persist through and be regulated by mitosis. Such nonhomologous interactions range from those ensuing from the clustering of PREs (GRIMAUD *et al.* 2006) and insulators scattered throughout the genome (reviewed by KUHN and GEYER 2003; CAPELSON and CORCES 2004; BRASSET and VAURY 2005; GASZNER and FELSENFELD 2006; VALENZUELA and KAMAKAKA 2006) to those occurring in transcription factories (reviewed by DILLON 2006; FARO-TRINDADE and COOK 2006; FRASER 2006). Excitingly, they also include gene-specific interactions, such as those that occur in mammals between enhancers and distant promoters during olfactory receptor choice (LOMVARDAS *et al.* 2006), between the interferon- γ and T_H2 cytokine loci during T-cell development (SPILIANAKIS *et al.* 2005), and between the insulin-like growth factor 2/H19 imprinting control region and the chromosomal segment containing the Wsb1 and Nf1 loci in a manner that appears to require the CCCTC binding factor CTCF (LING *et al.* 2006).

In closing, we note that while our studies have sought conditions that reduce somatic pairing in *Drosophila* cell culture, it will also be informative to seek situations in which pairing can be enhanced or induced. In particular, might the capacity to pair be an inherent property of chromosomes that is, however, precluded or suppressed in most organisms? Extending a suggestion by McClintock in 1933 of a “powerful force” that promotes pairing in meiosis (McCLINTOCK 1933), it may be that the paired state of chromosomes or chromosomal regions, be it homologous or not, is energetically, kinetically, or structurally favored (also see LEE *et al.* 2004) and must, therefore, be actively controlled. That somatic pairing may be a widespread potential of chromosomes is consistent with the abundance of homology- and pairing-dependent regulatory mechanisms as well as reports of nonhomologous interactions, while a necessity to suppress pairing is conceivable in situations where continual interallelic communication would be detrimental, such as may be the case when alleles are differentially regulated.

We thank Kami Ahmad for alerting us to the article by HALFER and BARIGOZZI (1973); Monica Colaiácovo for critical input and generous access to her DeltaVision Imaging System; the *Drosophila* Genomics Research Center for cell lines; Kami Ahmad, Frédéric Bantignies, Giacomo Cavalli, and George Church for technical advice and discussion; John Aach, Mar Carmena, Lucy Cherbas, Charleston Chiang, Stephen Elledge, Richard Emmons, Allan Gurtan, Scott Hawley, Amber Hohl, Tao Hsieh, Matt Jakubik, Mitzi Kuroda, Erica Larschan, Anne Lee, Jeannie Lee, Lillian Merriam, Norbert Perrimon, and Johannes Walter for thoughtful input; and Anna Moran and Laura Stadelmann for technical assistance. We apologize to authors we could not cite due to space limitations. This study was supported by a National Science Foundation Graduate Research Fellowship to B.R.W. and by a Ruth L. Kirschstein National Research Service Award (GM-67460) and a grant (GM-61936) from the National Institutes of Health to J.R.B. and C.-t.W., respectively.

LITERATURE CITED

- ANDOH, T., and R. ISHIDA, 1998 Catalytic inhibitors of DNA topoisomerase II. *Biochim. Biophys. Acta* **1400**: 155–171.
- AZUARA, V., K. E. BROWN, R. R. WILLIAMS, N. WEBB, N. DILLON *et al.*, 2003 Heritable gene silencing in lymphocytes delays chromatid resolution without affecting the timing of DNA replication. *Nat. Cell Biol.* **5**: 668–674.
- BACHANT, J., A. ALCASABAS, Y. BLAT, N. KLECKNER and S. J. ELLEDGE, 2002 The SUMO-1 isopeptidase Smt4 is linked to centromeric cohesion through SUMO-1 modification of DNA topoisomerase II. *Mol. Cell* **9**: 1169–1182.
- BACHER, C. P., M. GUGGIARI, B. BRORS, S. AUGUI, P. CLERC *et al.*, 2006 Transient colocalization of X-inactivation centres accompanies the initiation of X inactivation. *Nat. Cell Biol.* **8**: 293–299.
- BANTIGNIES, F., C. GRIMAUD, S. LAVROV, M. GABUT and G. CAVALLI, 2003 Inheritance of Polycomb-dependent chromosomal interactions in *Drosophila*. *Genes Dev.* **17**: 2406–2420.
- BANTIGNIES, F., C. GRIMAUD and G. CAVALLI, 2005 Two-colour fluorescent in situ DNA hybridization on whole mount *Drosophila* embryos and larval imaginal discs (PROT07). (<http://www.epigenome-noe.net/researchtools/protocol.php?protid=5>).
- BRASSET, E., and C. VAURY, 2005 Insulators are fundamental components of the eukaryotic genomes. *Heredity* **94**: 571–576.
- CAIRNS, J., 2006 Cancer and the immortal strand hypothesis. *Genetics* **174**: 1069–1072.
- CAPELSON, M., and V. G. CORCES, 2004 Boundary elements and nuclear organization. *Biol. Cell* **96**: 617–629.
- CHANG, C. J., S. GOULDING, W. C. EARNSHAW and M. CARMENA, 2003 RNAi analysis reveals an unexpected role for topoisomerase II in chromosome arm congression to a metaphase plate. *J. Cell Sci.* **116**: 4715–4726.
- CHEN, J. D., and V. PIRROTTA, 1993 Stepwise assembly of hyperaggregated forms of *Drosophila* zeste mutant protein suppresses white gene expression in vivo. *EMBO J.* **12**: 2061–2073.
- CHERBAS, L., 2007 *Drosophila* Genomics Resource Center (DGRC) protocols. (<http://dgrc.cgb.indiana.edu/cells/protocols.html>).
- CHERBAS, L., and P. CHERBAS, 1997 “Parahomologous” gene targeting in *Drosophila* cells: an efficient, homology-dependent pathway of illegitimate recombination near a target site. *Genetics* **145**: 349–358.
- CLEMENS, J. C., C. A. WORBY, N. SIMONSON-LEFF, M. MUDA, T. MAEHOMA *et al.*, 2000 Use of double-stranded RNA interference in *Drosophila* cell lines to dissect signal transduction pathways. *Proc. Natl. Acad. Sci. USA* **97**: 6499–6503.
- COBB, J., R. K. REDDY, C. PARK and M. A. HANDEL, 1997 Analysis of expression and function of topoisomerase I and II during meiosis in male mice. *Mol. Reprod. Dev.* **46**: 489–498.
- COELHO, P. A., J. QUEIROZ-MACHADO and C. E. SUNKEL, 2003 Condensin-dependent localisation of topoisomerase II to an axial chromosomal structure is required for sister chromatid resolution during mitosis. *J. Cell Sci.* **116**: 4763–4776.
- COLAIACOVO, M. P., 2006 The many facets of SC function during *C. elegans* meiosis. *Chromosoma* **115**: 195–211.
- CONBOY, M. J., A. O. KARASOV and T. A. RANDO, 2007 High incidence of non-random template strand segregation and asymmetric fate

- determination in dividing stem cells and their progeny. *PLoS Biol.* **5**: e102.
- CSINK, A. K., and S. HENIKOFF, 1998 Large-scale chromosomal movements during interphase progression in *Drosophila*. *J. Cell Biol.* **143**: 13–22.
- DEBEC, A., 1978 Haploid cell cultures of *Drosophila melanogaster*. *Nature* **274**: 255–256.
- DEBEC, A., 1984 Evolution of karyotype in haploid cell lines of *Drosophila melanogaster*. *Exp. Cell Res.* **151**: 236–246.
- DEJ, K. J., and A. C. SPRADLING, 1999 The endocycle controls nurse cell polytene chromosome structure during *Drosophila* oogenesis. *Development* **126**: 293–303.
- DERNBURG, A., 2000 *In situ* hybridization to somatic chromosomes, pp. 24–55 in *Drosophila Protocols*, edited by W. SULLIVAN, M. ASHBURNER and R. S. HAWLEY. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- DERNBURG, A. F., and J. W. SEDAT, 1998 Mapping three-dimensional chromosome architecture *in situ*. *Methods Cell Biol.* **53**: 187–233.
- DIAZ-PEREZ, S. V., D. O. FERGUSON, C. WANG, G. CSANKOVSKI, C. WANG *et al.*, 2006 A deletion at the mouse Xist gene exposes trans-effects that alter the heterochromatin of the inactive X chromosome and the replication time and DNA stability of both X chromosomes. *Genetics* **174**: 1115–1133.
- DILLON, N., 2006 Gene regulation and large-scale chromatin organization in the nucleus. *Chromosome Res.* **14**: 117–126.
- DONOHUE, M. E., L. F. ZHANG, N. XU, Y. SHI and J. T. LEE, 2007 Identification of a Ctcf cofactor, Yy1, for the X chromosome binary switch. *Mol. Cell* **25**: 43–56.
- DORSETT, D., 2007 Roles of the sister chromatid cohesion apparatus in gene expression, development, and human syndromes. *Chromosoma* **116**: 1–13.
- DUNCAN, I. W., 2002 Transvection effects in *Drosophila*. *Annu. Rev. Genet.* **36**: 521–556.
- EARNSHAW, W. C., and M. M. HECK, 1985 Localization of topoisomerase II in mitotic chromosomes. *J. Cell Biol.* **100**: 1716–1725.
- EARNSHAW, W. C., B. HALLIGAN, C. A. COOKE, M. M. HECK and L. F. LIU, 1985 Topoisomerase II is a structural component of mitotic chromosome scaffolds. *J. Cell Biol.* **100**: 1706–1715.
- ECHALIER, G., 1997 *Drosophila Cells in Culture*. Academic Press, San Diego.
- ECHALIER, G., and A. OHANESSIAN, 1969 Isolation, in tissue culture, of *Drosophila melanogaster* cell lines. *C. R. Acad. Sci. Hebd. Seances Acad. Sci. D* **268**: 1771–1773.
- FARO-TRINDADE, I., and P. R. COOK, 2006 Transcription factories: structures conserved during differentiation and evolution. *Biochem. Soc. Trans.* **34**: 1133–1137.
- FRASER, P., 2006 Transcriptional control thrown for a loop. *Curr. Opin. Genet. Dev.* **16**: 490–495.
- FRITSCH, C., G. PLOEGER and D. J. ARNDT-JOVIN, 2006 *Drosophila* under the lens: imaging from chromosomes to whole embryos. *Chromosome Res.* **14**: 451–464.
- FUNG, J. C., W. F. MARSHALL, A. DERNBURG, D. A. AGARD and J. W. SEDAT, 1998 Homologous chromosome pairing in *Drosophila melanogaster* proceeds through multiple independent initiations. *J. Cell Biol.* **141**: 5–20.
- GASSER, S. M., T. LAROCHE, J. FALQUET, E. BOY DE LA TOUR and U. K. LAEMMLI, 1986 Metaphase chromosome structure. Involvement of topoisomerase II. *J. Mol. Biol.* **188**: 613–629.
- GASZNER, M., and G. FELSENFELD, 2006 Insulators: exploiting transcriptional and epigenetic mechanisms. *Nat. Rev. Genet.* **7**: 703–713.
- GEMKOW, M. J., P. J. VERVEER and D. J. ARNDT-JOVIN, 1998 Homologous association of the Bithorax-Complex during embryogenesis: consequences for transvection in *Drosophila melanogaster*. *Development* **125**: 4541–4552.
- GERTON, J. L., and R. S. HAWLEY, 2005 Homologous chromosome interactions in meiosis: diversity amidst conservation. *Nat. Rev. Genet.* **6**: 477–487.
- GLOOR, G. B., 2002 The role of sequence homology in the repair of DNA double-strand breaks in *Drosophila*. *Adv. Genet.* **46**: 91–117.
- GOLIC, M. M., and K. G. GOLIC, 1996 A quantitative measure of the mitotic pairing of alleles in *Drosophila melanogaster* and the influence of structural heterozygosity. *Genetics* **143**: 385–400.
- GRANT-DOWNTON, R. T., and H. G. DICKINSON, 2004 Plants, pairing and phenotypes—Two's company? *Trends Genet.* **20**: 188–195.
- GRIMAUD, C., F. BANTIGNIES, M. PAL-BHADRA, P. GHANA, U. BHADRA *et al.*, 2006 RNAi components are required for nuclear clustering of Polycomb group response elements. *Cell* **124**: 957–971.
- GUBB, D., J. ROOTE, J. TRENEAR, D. COULSON, and M. ASHBURNER, 1997 Topological constraints on transvection between white genes within the transposing element TE35B in *Drosophila melanogaster*. *Genetics* **146**: 919–937.
- HAARS, R., H. ZENTGRAF, E. GATEFF and F. A. BAUTZ, 1980 Evidence for endogenous reovirus-like particles in a tissue culture cell line from *Drosophila melanogaster*. *Virology* **101**: 124–130.
- HALFER, C., and C. BARIGOZZI, 1973 Prophase synapsis in somatic cells of *Drosophila melanogaster*. *Chromosomes Today* **4**: 181–186.
- HOLT, C. M., 1917 Multiple complexes in the alimentary tract of *Culex pipiens*. *J. Morphol.* **29**: 607–627.
- HOSSAIN, M. S., K. KUROKAWA, N. AKIMITSU and K. SEKIMIZU, 2004 DNA topoisomerase II is required for the G0-to-S phase transition in *Drosophila* Schneider cells, but not in yeast. *Genes Cells* **9**: 905–917.
- IWABATA, K., A. KOSHIYAMA, T. YAMAGUCHI, H. SUGAWARA, F. N. HAMADA *et al.*, 2005 DNA topoisomerase II interacts with Lim15/Dmcl in meiosis. *Nucleic Acids Res.* **33**: 5809–5818.
- JU, B. G., V. V. LUNYAK, V. PERISSI, I. GARCIA-BASSETS, D. W. ROSE *et al.*, 2006 A topoisomerase IIbeta-mediated dsDNA break required for regulated transcription. *Science* **312**: 1798–1802.
- KASSIS, J. A., 2002 Pairing-sensitive silencing, polycomb group response elements, and transposon homing in *Drosophila*. *Adv. Genet.* **46**: 421–438.
- KAVI, H. H., H. R. FERNANDEZ, W. XIE and J. A. BIRCHLER, 2006 Polycomb, pairing and PIWI-RNA silencing and nuclear interactions. *Trends Biochem. Sci.* **31**: 485–487.
- KENNISON, J. A., and J. W. SOUTHWORTH, 2002 Transvection in *Drosophila*. *Adv. Genet.* **46**: 399–420.
- KOPCZYNSKI, C. C., and M. A. MUSKAVITCH, 1992 Introns excised from the Delta primary transcript are localized near sites of Delta transcription. *J. Cell Biol.* **119**: 503–512.
- KRAVCHENKO, E., E. SAVITSKAYA, O. KRAVCHUK, A. PARSHIKOV, P. GEORGIEV *et al.*, 2005 Pairing between gypsy insulators facilitates the enhancer action in trans throughout the *Drosophila* genome. *Mol. Cell Biol.* **25**: 9283–9291.
- KUHN, E. J., and P. K. GEYER, 2003 Genomic insulators: connecting properties to mechanism. *Curr. Opin. Cell Biol.* **15**: 259–265.
- LARSEN, A. K., A. E. ESCARGUEIL and A. SKLADANOWSKI, 2003 Catalytic topoisomerase II inhibitors in cancer therapy. *Pharmacol. Ther.* **99**: 167–181.
- LASALLE, J. M., and M. LALANDE, 1996 Homologous association of oppositely imprinted chromosomal domains. *Science* **272**: 725–728.
- LEE, D. J., A. WYNVEEN and A. A. KORNYUSHEV, 2004 DNA-DNA interaction beyond the ground state. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **70**: 051913.
- LING, J. Q., T. LI, J. F. HU, T. H. VU, H. L. CHEN *et al.*, 2006 CTCF mediates interchromosomal colocalization between Igf2/H19 and Wsb1/Nfl. *Science* **312**: 269–272.
- LOMVARDAS, S., G. BARNEA, D. J. PISAPIA, M. MENDELSON, J. KIRKLAND *et al.*, 2006 Interchromosomal interactions and olfactory receptor choice. *Cell* **126**: 403–413.
- LUPO, R., A. BREILING, M. E. BIANCHI and V. ORLANDO, 2001 *Drosophila* chromosome condensation proteins Topoisomerase II and Barren colocalize with Polycomb and maintain Fab-7 PRE silencing. *Mol. Cell* **7**: 127–136.
- MAESHIMA, K., and U. K. LAEMMLI, 2003 A two-step scaffolding model for mitotic chromosome assembly. *Dev. Cell* **4**: 467–480.
- MAESHIMA, K., M. ELTSOV and U. K. LAEMMLI, 2005 Chromosome structure: improved immunolabeling for electron microscopy. *Chromosoma* **114**: 365–375.
- MARAHRENS, Y., 1999 X-inactivation by chromosomal pairing events. *Genes Dev.* **13**: 2624–2632.
- MARSHALL, W. F., A. F. DERNBURG, B. HARMON, D. A. AGARD and J. W. SEDAT, 1996 Specific interactions of chromatin with the nuclear envelope: positional determination within the nucleus in *Drosophila melanogaster*. *Mol. Biol. Cell* **7**: 825–842.
- MCCCLINTOCK, B., 1933 The association of non-homologous parts of chromosomes in the midprophase of meiosis in *Zea mays*. *Z. Zellforsch. Mikrok. Anat.* **19**: 191–237.
- MCKEE, B. D., 2004 Homologous pairing and chromosome dynamics in meiosis and mitosis. *Biochim. Biophys. Acta* **1677**: 165–180.

- METZ, C. W., 1916 Chromosome studies on the Diptera. II. The paired association of chromosomes in the Diptera and its significance. *J. Exp. Zool.* **21**: 213–279.
- METZ, C. W., 1922 Association of homologous chromosomes in tetraploid cells of Diptera. *Biol. Bull.* **43**: 369–373.
- METZ, C. W., 1925 Prophase chromosome behavior in triploid individuals of *Drosophila melanogaster*. *Genetics* **10**: 345–350.
- MLYNARCZYK-EVANS, S., M. ROYCE-TOLLAND, M. K. ALEXANDER, A. A. ANDERSEN, S. KALANTRY *et al.*, 2006 X chromosomes alternate between two states prior to random X-inactivation. *PLoS Biol.* **4**: e159.
- NABIROCHKIN, S., M. OSSOKINA and T. HEIDMANN, 1998 A nuclear matrix/scaffold attachment region co-localizes with the gypsy retrotransposon insulator sequence. *J. Biol. Chem.* **273**: 2473–2479.
- NICHOLS, W. W., C. BRADT and W. BOWNE, 1971 V. Cytogenetic studies on cells in culture from the class Insecta. *Curr. Top. Microbiol. Immunol.* **55**: 61–69.
- OSMAN, K., E. SANCHEZ-MORAN, J. D. HIGGINS, G. H. JONES and F. C. FRANKLIN, 2006 Chromosome synapsis in Arabidopsis: analysis of the transverse filament protein ZYP1 reveals novel functions for the synaptonemal complex. *Chromosoma* **115**: 212–219.
- PAWLOWSKI, W. P., and W. Z. CANDE, 2005 Coordinating the events of the meiotic prophase. *Trends Cell Biol.* **15**: 674–681.
- PEEL, D. J., and M. J. MILNER, 1991 Karyotype analysis of imaginal disc cell lines. *Dros. Inf. Serv.* **70**: 176–177.
- PEEL, D. J., S. A. JOHNSON and M. J. MILNER, 1990 The ultrastructure of imaginal disc cells in primary cultures and during cell aggregation in continuous cell lines. *Tissue Cell* **22**: 749–758.
- PERRIMON, N., 2007 RNAi protocols. (http://flyrnai.org/RNAi_protocols.html).
- PIRROTTA, V., 1999 Transvection and chromosomal trans-interaction effects. *Biochim. Biophys. Acta* **1424**: M1–M8.
- POMMIER, Y., 2006 Topoisomerase I inhibitors: camptothecins and beyond. *Nat. Rev. Cancer* **6**: 789–802.
- PORTER, A. C., and C. J. FARR, 2004 Topoisomerase II: untangling its contribution at the centromere. *Chromosome Res.* **12**: 569–583.
- RIESELTMANN, L., and T. HAAF, 1999 Preferential S-phase pairing of the imprinted region on distal mouse chromosome 7. *Cytogenet. Cell Genet.* **86**: 39–42.
- RONG, Y. S., and K. G. GOLIC, 2003 The homologous chromosome is an effective template for the repair of mitotic DNA double-strand breaks in *Drosophila*. *Genetics* **165**: 1831–1842.
- RONSHAUGEN, M., and M. LEVINE, 2004 Visualization of trans-homolog enhancer-promoter interactions at the Abd-B Hox locus in the *Drosophila* embryo. *Dev. Cell* **7**: 925–932.
- SASS, G. L., and S. HENIKOFF, 1999 Pairing-dependent mislocalization of a *Drosophila* brown gene reporter to a heterochromatic environment. *Genetics* **152**: 595–604.
- SAVVIDOU, E., N. COBBE, S. STEFFENSEN, S. COTTERILL and M. M. HECK, 2005 *Drosophila* CAP-D2 is required for condensin complex stability and resolution of sister chromatids. *J. Cell Sci.* **118**: 2529–2543.
- SCHNEIDER, L., 1972 Cell lines derived from late embryonic stages of *Drosophila melanogaster*. *J. Embryol. Exp. Morphol.* **27**: 353–365.
- SILAHTAROGLU, A., H. PFUNDHELLER, A. KOSHKIN, N. TOMMERUP and S. KAUPPINEN, 2004 LNA-modified oligonucleotides are highly efficient as FISH probes. *Cytogenet. Genome Res.* **107**: 32–37.
- SILAHTAROGLU, A. N., N. TOMMERUP and H. VISSING, 2003 FISHing with locked nucleic acids (LNA): evaluation of different LNA/DNA mixers. *Mol. Cell. Probes* **17**: 165–169.
- SMILEY, R. D., T. R. COLLINS, G. G. HAMMES and T. S. HSIEH, 2007 Single-molecule measurements of the opening and closing of the DNA gate by eukaryotic topoisomerase II. *Proc. Natl. Acad. Sci. USA* **104**: 4840–4845.
- SOLIER, S., A. LANSIAUX, E. LOGETTE, J. WU, J. SORET *et al.*, 2004 Topoisomerase I and II inhibitors control caspase-2 pre-messenger RNA splicing in human cells. *Mol. Cancer Res.* **2**: 53–61.
- SONDERMEIJER, P. J., J. W. DERKSEN and N. H. LUBSEN, 1980 New cell line: established cell lines of *Drosophila hydei*. *In Vitro* **16**: 913–914.
- SPILIANAKIS, C. G., M. D. LALIOTI, T. TOWN, G. R. LEE and R. A. FLAVELL, 2005 Interchromosomal associations between alternatively expressed loci. *Nature* **435**: 637–645.
- STEVENS, N., 1907 The chromosomes in *Drosophila ampelophila*. Proceedings of the 7th International Zoological Congress. Cambridge University Press, Cambridge, UK.
- STEVENS, N., 1908 A study of the germ cells of certain Diptera, with reference to the heterochromosomes and phenomena of synapsis. *J. Exp. Zool.* **5**: 359–374.
- THATCHER, K. N., S. PEDDADA, D. H. YASUI and J. M. LASALLE, 2005 Homologous pairing of 15q11–13 imprinted domains in brain is developmentally regulated but deficient in Rett and autism samples. *Mol. Genet.* **14**: 785–797.
- TSUBOUCHI, T., and G. S. ROEDER, 2005 A synaptonemal complex protein promotes homology-independent centromere coupling. *Science* **308**: 870–873.
- UI, K., S. NISHIHARA, M. SAKUMA, S. TOGASHI, R. UEDA *et al.*, 1994 Newly established cell lines from *Drosophila* larval CNS express neural specific characteristics. *In Vitro Cell Dev. Biol. Anim.* **30A**: 209–216.
- VALENZUELA, L., and R. T. KAMAKAKA, 2006 Chromatin insulators. *Annu. Rev. Genet.* **40**: 107–138.
- VAN HOOSER, A. A., P. YUH and R. HEALD, 2005 The perichromosomal layer. *Chromosoma* **114**: 377–388.
- VAZQUEZ, J., A. S. BELMONT and J. W. SEDAT, 2002 The dynamics of homologous chromosome pairing during male *Drosophila* meiosis. *Curr. Biol.* **12**: 1473–1483.
- VAZQUEZ, J., M. MULLER, V. PIRROTTA and J. W. SEDAT, 2006 The Mcp element mediates stable long-range chromosome-chromosome interactions in *Drosophila*. *Mol. Biol. Cell* **17**: 2158–2165.
- VEILE, R., 1990 Chromosome analysis methods (http://hg.wustl.edu/hdk_lab_manual/cam/camcontents.html).
- WANG, X., and W. DAI, 2005 Shugoshin, a guardian for sister chromatid segregation. *Exp. Cell Res.* **310**: 1–9.
- WATANABE, Y., 2005 Sister chromatid cohesion along arms and at centromeres. *Trends Genet.* **21**: 405–412.
- WU, C. T., and J. R. MORRIS, 1999 Transvection and other homology effects. *Curr. Opin. Genet. Dev.* **9**: 237–246.
- WYMAN, C., D. RISTIC and R. KANAAR, 2004 Homologous recombination-mediated double-strand break repair. *DNA Repair* **3**: 827–833.
- XU, N., C. L. TSAI and J. T. LEE, 2006 Transient homologous chromosome pairing marks the onset of X inactivation. *Science* **311**: 1107–1109.
- YANAGAWA, S., J. S. LEE and A. ISHIMOTO, 1998 Identification and characterization of a novel line of *Drosophila* Schneider S2 cells that respond to wingless signaling. *J. Biol. Chem.* **273**: 32353–32359.
- ZICKLER, D., 2006 From early homologue recognition to synaptonemal complex formation. *Chromosoma* **115**: 158–174.

Communicating editor: J. A. BIRCHLER